

CRITICAL POINTS IN AN ALGEBRA OF ELEMENTARY EMBEDDINGS

RANDALL DOUGHERTY

Ohio State University

May 27, 1993

ABSTRACT. Given two elementary embeddings from the collection of sets of rank less than λ to itself, one can combine them to obtain another such embedding in two ways: by composition, and by applying one to (initial segments of) the other. Hence, a single such nontrivial embedding j generates an algebra of embeddings via these two operations, which satisfies certain laws (for example, application distributes over both composition and application). Laver has shown, among other things, that this algebra is free on one generator with respect to these laws.

The set of critical points of members of this algebra is the subject of this paper. This set contains the critical point κ_0 of j , as well as all of the other ordinals κ_n in the critical sequence of j (defined by $\kappa_{n+1} = j(\kappa_n)$). But the set includes many other ordinals as well. The main result of this paper is that the number of critical points below κ_n (which has been shown to be finite by Laver and Steel) grows so quickly with n that it dominates any primitive recursive function. In fact, it grows faster than the Ackermann function, and even faster than a slow iterate of the Ackermann function. Further results show that, even just below κ_4 , one can find so many critical points that the number is only expressible using fast-growing hierarchies of iterated functions (six levels of iteration beyond exponentials).

1. INTRODUCTION

Let V_λ be the collection of sets of rank less than λ , where λ is some fixed limit ordinal. The assumption that there is a nontrivial elementary embedding from V_λ to V_λ is an extremely strong large cardinal hypothesis [8]. But once one such embedding is known to exist, more of them can be obtained by applying embeddings to each other. If a and b are two such embeddings, then one cannot literally apply a to b since $b \notin V_\lambda$, but one can apply a to initial segments of b , so we define $a(b)$ to be $\bigcup_{\alpha < \lambda} a(b \upharpoonright V_\alpha)$. Then $a(b)$ will also be an elementary embedding from V_λ to V_λ . Of course, one can also obtain new embeddings from old ones by composition.

Let $\mathcal{A}_j$ be the collection of embeddings generated by a single nontrivial embedding $j: V_\lambda \rightarrow V_\lambda$ using the application operation. Elementarity implies that application is left distributive over itself: $a(b(c)) = a(b)(a(c))$. Laver [5] has shown that $\mathcal{A}_j$ is a free algebra on the generator j with respect to the left distributive law. Dehornoy [2] has carried out an algebraic study of left distributivity; in particular, Dehornoy [1] proved a property (irreflexivity) of the free left distributive algebra on one generator which previously had only been known under the above large cardinal assumption [5]. (A simplification of the proof of this particular result has since been given by Larue [4].)

If one considers the larger algebra $\mathcal{P}_j$ generated by j using both application and composition, then one immediately gets the laws $a \circ (b \circ c) = (a \circ b) \circ c$ and $(a \circ b)(c) = a(b(c))$, and elementarity of

The author was supported by NSF grant number DMS-9158092.

a gives $a(b \circ c) = a(b) \circ a(c)$ and $a \circ b = a(b) \circ a$ (the latter because $a(b(x)) = a(b)(a(x))$). Laver [5] shows that $\mathcal{P}_j$ is free with respect to these four laws.

Any embedding $k \in \mathcal{P}_j$ maps ordinals to ordinals in a strictly increasing manner, so $k(\alpha) \geq \alpha$ for all $\alpha < \lambda$; the least α such that $k(\alpha) > \alpha$ is called the *critical point* of k , and denoted by $cr(k)$. This α is inaccessible, measurable, etc., and $k(x) = x$ for $x \in V_\alpha$ [8]. Elementarity implies that $k'(cr(k)) = cr(k'(k))$ and $k'(k(\beta)) = k'(k)(k'(\beta))$; using these rules, we can obtain many ordinals below λ as critical points of members of $\mathcal{A}_j$. To start with, let $\kappa_0 = cr(j)$ and $\kappa_{n+1} = j(\kappa_n)$ for $n \in \omega$; the ordinals κ_n form a strictly increasing sequence (called the *critical sequence* of j), and all of them are critical points of members of $\mathcal{A}_j$ (if $\kappa_n = cr(k)$, then $\kappa_{n+1} = cr(j(k))$). Perhaps surprisingly, it turns out that other ordinals also occur as critical points. To see this, define the sequence of embeddings $j_{(n)}$ by $j_{(1)} = j$ and $j_{(n+1)} = j_{(n)}(j)$; then we easily compute $cr(j_{(2)}) = j(\kappa_0) = \kappa_1$, $j_{(2)}(\kappa_1) = j(j)(j(\kappa_0)) = j(j(\kappa_0)) = \kappa_2$, $j_{(2)}(\kappa_2) = \kappa_3$, $cr(j_{(3)}) = j_{(2)}(\kappa_0) = \kappa_0$, $j_{(3)}(\kappa_0) = j_{(2)}(j)(j_{(2)}(\kappa_0)) = j_{(2)}(j(\kappa_0)) = \kappa_2$, and $j_{(3)}(\kappa_2) = \kappa_3$, so $j_{(3)}(\kappa_1) = cr(j_{(3)}(j_{(2)}))$ must lie strictly between κ_2 and κ_3 .

Further computations yield a number of critical points between κ_3 and κ_4 (for example, the ordinals $j(j_{(3)}(\kappa_1))$, $j_{(2)}(j_{(3)}(\kappa_1))$, and $j_{(3)}(j_{(3)}(\kappa_1))$ turn out to be distinct). In fact, one cannot immediately rule out the possibility that there are infinitely many critical points below κ_4 . This turns out not to be the case; results of Steel and Laver [6] show that the collection of critical points of members of $\mathcal{P}_j$ has order type ω . (To be precise, Laver showed that the ordinals $cr(j_{(n)})$, $n = 1, 2, \dots$ are the first ω critical points of members of $\mathcal{P}_j$, listed with duplications, and Steel showed that the ordinals $cr(j_{(n)})$, $n = 1, 2, \dots$ are cofinal in λ .)

The main result of this paper is that there are very many critical points other than the ordinals κ_n .

Theorem 1. *If $F(n)$ is the number of critical points of members of $\mathcal{P}_j$ lying below κ_n , then F grows faster than any primitive recursive function of n .*

This can be made more precise by considering the standard fast-growing hierarchy of functions. (There are slight variations in the “standard” definition of this hierarchy. The version used here is based on a formula of Hermes and Péter, which makes the calculations in this paper simpler; using another version would not change results like Theorem 1, but it would affect the explicit formulas relating the numbers of critical points to the hierarchy. See Rose [7] for further details.) Let $F_0(k) = k + 1$, and define F_{n+1} by iterating F_n : $F_{n+1}(0) = F_n(1)$, $F_{n+1}(k+1) = F_n(F_{n+1}(k))$. So $F_3(k) = 2^{k+3} - 3$, F_4 is an iterated exponential, and so on. Then diagonalize to obtain $F_\omega(k) = F_k(k)$, and iterate F_ω to get $F_{\omega+1}$, etc. Standard results show that any primitive recursive function is eventually dominated by F_n for some finite n , and therefore by F_ω (which is a variant of the Ackermann function).

An easy induction on terms in $\mathcal{P}_j$ shows that any element of $\mathcal{P}_j$ can be written as a composition of one or more members of $\mathcal{A}_j$. Since the critical point of a composition of embeddings is the minimum of the critical points of the individual embeddings, this shows that all critical points of members of $\mathcal{P}_j$ are actually critical points of members of $\mathcal{A}_j$. (In fact, except in section 3, all of the embeddings used in the constructions are actually in $\mathcal{A}_j$.)

Section 2 of this paper is a proof that $F(n) > F_\omega(n-1)$ for all $n \geq 4$; this suffices to prove the above theorem. In section 3 this result is improved to: $F(n) > F_{\omega+1}(\lfloor \log_3 n \rfloor - 1)$ for all $n \geq 3$. These results are strong asymptotically, but do not say much about $F(n)$ for specific small n . It turns out that $F(3) = 4$, but $F(4)$ is quite large; section 4 gives most of a proof that $F(4)$ exceeds $F_9(F_8(F_8(254)))$, which is an incomprehensibly large number. Section 5 covers computation of critical point inequalities, which leads to some interesting problems; this section can

be read independently of sections 2–4, although one of its purposes is to provide a few inequalities needed to complete the proof in section 4.

It follows from the theorem above that the algebraic content of the Steel-Laver result on critical points (see Laver [6] and Dougherty and Jech [3]) cannot be proved by primitive recursive methods (i.e., cannot be proved in the standard theory of Primitive Recursive Arithmetic). This contrasts with the irreflexivity property of the free algebra, which, as Dehornoy shows, is provable by primitive recursive methods.

2. THE MAIN CONSTRUCTION

We will now prove Theorem 1 through a sequence of lemmas. The lemmas will be self-contained, but the full construction given by them may be hard to visualize; therefore, after the proof is complete, a picture of the whole construction and some additional explanatory comments will be given.

Throughout the rest of the paper, ‘critical point’ will mean ‘critical point of a member of $\mathcal{P}_j$ ’ (which, as shown in the preceding section, is the same as ‘critical point of a member of $\mathcal{A}_j$ ’), and ‘embedding’ will mean ‘member of $\mathcal{P}_j$.’

We start by computing the critical sequences of certain simple members of $\mathcal{A}_j$. By definition, the critical sequence of j itself is $\langle \kappa_n : n \in \omega \rangle$, or, in a more suggestive notation,

$$\kappa_0 \mapsto \kappa_1 \mapsto \kappa_2 \mapsto \kappa_3 \mapsto \cdots .$$

Also, if the critical sequence of i is

$$\mu_0 \mapsto \mu_1 \mapsto \mu_2 \mapsto \mu_3 \mapsto \cdots ,$$

then the critical sequence of $k(i)$ is

$$k(\mu_0) \mapsto k(\mu_1) \mapsto k(\mu_2) \mapsto k(\mu_3) \mapsto \cdots .$$

Therefore, if we define the sequence of embeddings $j^{[n]}$ by $j^{[0]} = j$ and $j^{[n+1]} = j(j^{[n]})$, then $j^{[n]}$ has critical sequence

$$\kappa_n \mapsto \kappa_{n+1} \mapsto \kappa_{n+2} \mapsto \kappa_{n+3} \mapsto \cdots .$$

By applying the embeddings $j^{[n]}$ to each other, we can get embeddings with slightly more complicated critical sequences. For example, the embedding $j^{[1]}(j)$ has critical sequence

$$\kappa_0 \mapsto \kappa_2 \mapsto \kappa_3 \mapsto \kappa_4 \mapsto \cdots ,$$

and the embedding $j^{[3]}(j^{[3]}(j^{[1]}))$ has critical sequence

$$\kappa_1 \mapsto \kappa_2 \mapsto \kappa_5 \mapsto \kappa_6 \mapsto \kappa_7 \mapsto \cdots .$$

In general, given any finite specification

$$\kappa_{n(0)} \mapsto \kappa_{n(1)} \mapsto \cdots \mapsto \kappa_{n(l)}$$

with $n(0) < n(1) < \cdots < n(l)$, one can find a member of $\mathcal{A}_j$ whose critical sequence begins with the given specification: start with j , apply j $n(0)$ times to move the critical point up to $\kappa_{n(0)}$,

then apply $j^{[n(0)+1]}$ $n(1) - n(0) - 1$ times to move the next member of the critical sequence up to $\kappa_{n(1)}$, then apply $j^{[n(1)+1]}$ $n(2) - n(1) - 1$ times, and so on. The examples below will only use some simple cases of this; for instance, if one wants an embedding k with critical sequence beginning $\kappa_2 \mapsto \kappa_n \mapsto \kappa_{n+1}$ ($n \geq 3$), then one can let $k = (j^{[3]})^{n-3}(j^{[2]})$ (i.e., start with $j^{[2]}$ and apply $j^{[3]}$ $n - 3$ times).

A main fact to be used in the construction is that each embedding in $\mathcal{P}_j$ is a strictly order-preserving map from critical points to critical points; this will be used over and over again to produce more and more critical points, and we will similarly get more and more embeddings as well. To start with:

Lemma 2. *Suppose that one has critical points $\zeta < \alpha_0 < \alpha_1 < \dots < \alpha_n$ and embeddings $a_i \in \mathcal{P}_j$ for $i < n$ such that a_i has critical sequence beginning $\zeta \mapsto \alpha_i \mapsto \alpha_{i+1}$ for each i . Then there are at least 2^n critical points in the half-open interval $[\zeta, \alpha_n)$.*

Proof. Induct on n ; for $n = 0$, the single critical point ζ will suffice. Given the result for n , if one has α_i for $i \leq n + 1$ and a_i for $i < n + 1$ as above, then the induction hypothesis gives at least 2^n critical points in $[\zeta, \alpha_n)$. The embedding a_n maps the interval $[\zeta, \alpha_n)$ to the interval $[\alpha_n, \alpha_{n+1})$, and maps distinct critical points in the former interval to distinct critical points in the latter. Therefore, there are at least 2^n critical points in $[\alpha_n, \alpha_{n+1})$, giving a total of at least 2^{n+1} critical points in $[\zeta, \alpha_{n+1})$. This completes the induction. ■

For example, if we let $\zeta = \kappa_0$, $\alpha_i = \kappa_{i+1}$, and $a_i = (j^{[1]})^i(j)$, then we get at least 2^n critical points below κ_{n+1} for any n . However, we can do much better than this.

Lemma 3. *Suppose that one has critical points $\zeta < \zeta' < \beta_0 < \beta_1 < \dots < \beta_n$ and embeddings $b_i \in \mathcal{P}_j$ for $i < n$ such that b_i has critical sequence beginning $\zeta' \mapsto \beta_i \mapsto \beta_{i+1}$ for each i . Also suppose that there is an embedding z such that z has critical sequence beginning $\zeta \mapsto \zeta' \mapsto \beta_0$. Then there exist critical points α_i ($i \leq 2^n$) and embeddings a_i ($i < 2^n$) such that a_i has critical sequence beginning $\zeta \mapsto \alpha_i \mapsto \alpha_{i+1}$ for each i , $\alpha_0 = \zeta'$, and $\alpha_{2^n} = \beta_n$.*

Proof. Induct on n ; for $n = 0$, just let $a_0 = z$, $\alpha_0 = \zeta'$, and $\alpha_1 = \beta_0$. Given the result for n , if one has β_i for $i \leq n + 1$ and b_i for $i < n + 1$ as above, then the induction hypothesis gives critical points α_i ($i \leq 2^n$) and embeddings a_i ($i < 2^n$) as required, with $\alpha_0 = \zeta'$ and $\alpha_{2^n} = \beta_n$. Define α_{2^n+i} for $1 \leq i \leq 2^n$ by the equation $\alpha_{2^n+i} = b_n(\alpha_i)$; note that this equation also holds for $i = 0$, since $b_n(\zeta') = \beta_n$. Also note that $b_n(\zeta) = \zeta$, since $\zeta < \zeta' = cr(b_n)$. Hence, if we let $a_{2^n+i} = b_n(a_i)$ for $0 \leq i < 2^n$, then a_{2^n+i} has critical sequence beginning $\zeta \mapsto \alpha_{2^n+i} \mapsto \alpha_{2^n+i+1}$ for each such i . Finally, $\alpha_{2^{n+1}} = b_n(\beta_n) = \beta_{n+1}$. This completes the induction. ■

Note that it follows from the fact that a_i has critical sequence beginning $\zeta \mapsto \alpha_i \mapsto \alpha_{i+1}$ that $\zeta < \alpha_i < \alpha_{i+1}$.

One could apply this lemma with $\zeta = \kappa_0$, $\zeta' = \kappa_1$, $\beta_i = \kappa_{i+2}$, $z = j$, and $b_i = (j^{[2]})^i(j^{[1]})$ to get 2^n embeddings as in Lemma 2 ending up with $\alpha_{2^n} = \kappa_{n+2}$; hence, Lemma 2 would give 2^{2^n} critical points below κ_{n+2} . In fact, for any m , if one started with embeddings b_i ($i \leq n$) such that b_i has critical sequence beginning $\kappa_m \mapsto \kappa_{m+1+i} \mapsto \kappa_{m+2+i}$, then one could apply Lemma 3 m times and then apply Lemma 2 to get $\exp_2^{m+1}(n)$ (a tower of exponentials consisting of $m + 1$ 2's with an n at the top) critical points below κ_{n+m+1} . Again, however, this is not as much as following results will give.

The next result uses a numerical function $C_2(n)$ defined recursively as follows: $C_2(0) = 0$ and $C_2(n + 1) = C_2(n) + 2^{C_2(n)}$. (So C_2 is something like an iterated exponential.)

Lemma 4. *Suppose that one has critical points $\zeta < \zeta' < \delta_0 < \delta_1 < \dots < \delta_n$ and embeddings $d_i \in \mathcal{P}_j$ for $i < n$ such that d_i has critical sequence beginning $\zeta \mapsto \zeta' \mapsto \delta_i \mapsto \delta_{i+1}$ for each i . Then there exist critical points β_i ($i \leq C_2(n)$) and embeddings b_i ($i < C_2(n)$) such that b_i has critical sequence beginning $\zeta' \mapsto \beta_i \mapsto \beta_{i+1}$ for each i , $\beta_0 = \delta_0$, and $\beta_{C_2(n)} = \delta_n$.*

Proof. Induct on n ; the case $n = 0$ is essentially vacuous (just let $\beta_0 = \delta_0$). Given the result for n , if one has δ_i for $i \leq n + 1$ and d_i for $i < n + 1$ as above, then the induction hypothesis gives critical points β_i ($i \leq C_2(n)$) and embeddings b_i ($i < C_2(n)$) as required, with $\beta_0 = \delta_0$ and $\beta_{C_2(n)} = \delta_n$. We can now apply Lemma 3, using $z = d_0$, to get critical points α_i ($i \leq m$) and embeddings a_i ($i < m$), where $m = 2^{C_2(n)}$, such that a_i has critical sequence beginning $\zeta \mapsto \alpha_i \mapsto \alpha_{i+1}$ for each i , $\alpha_0 = \zeta'$, and $\alpha_m = \beta_{C_2(n)} = \delta_n$. Define $\beta_{C_2(n)+i}$ for $1 \leq i \leq m$ by the equation $\beta_{C_2(n)+i} = d_n(\alpha_i)$; note that this equation also holds for $i = 0$, since $d_n(\zeta') = \delta_n$. Hence, if we let $b_{C_2(n)+i} = d_n(a_i)$ for $0 \leq i < m$, then $b_{C_2(n)+i}$ has critical sequence beginning $\zeta' \mapsto \beta_{C_2(n)+i} \mapsto \beta_{C_2(n)+i+1}$ for each such i . Finally, $\beta_{C_2(n)+1} = d_n(\delta_n) = \delta_{n+1}$. This completes the induction. ■

Lemma 5. *Suppose that one has critical points $\zeta < \zeta' < \zeta'' < \varepsilon_0 < \varepsilon_1 < \dots < \varepsilon_n$ and embeddings $e_i \in \mathcal{P}_j$ for $i < n$ such that e_i has critical sequence beginning $\zeta'' \mapsto \varepsilon_i \mapsto \varepsilon_{i+1}$ for each i . Also suppose that there is an embedding z such that z has critical sequence beginning $\zeta \mapsto \zeta' \mapsto \zeta'' \mapsto \varepsilon_0$. Then there exist critical points δ_i ($i \leq 2^n$) and embeddings d_i ($i < 2^n$) such that d_i has critical sequence beginning $\zeta \mapsto \zeta' \mapsto \delta_i \mapsto \delta_{i+1}$ for each i , $\delta_0 = \zeta''$, and $\delta_{2^n} = \varepsilon_n$.*

Proof. Just as in the proof of Lemma 3, let $d_0 = z$ and $d_{2^n+i} = e_n(d_i)$ for $0 \leq i < 2^n$. ■

We need one more lemma of this sort, for which we need a family of numerical functions to be denoted $C_{2N+2}(n)$ for $N \geq 1$. These are again defined recursively, both on N and n , by the formulas $C_{2N+2}(0) = 0$ and $C_{2N+2}(n+1) = C_{2N+2}(n) + C_{2N}(2^{C_{2N+2}(n)})$. (So C_{2N+2} is an augmented iteration of C_{2N} .)

Lemma 6. *Let $N \geq 1$, and suppose that one has critical points $\zeta_0 < \zeta_1 < \dots < \zeta_{N+1} < \eta_0 < \eta_1 < \dots < \eta_n$ and embeddings $h_i \in \mathcal{P}_j$ for $i < n$ such that h_i has critical sequence beginning $\zeta_N \mapsto \zeta_{N+1} \mapsto \eta_i \mapsto \eta_{i+1}$ for each i . Also suppose that there exist embeddings z_k ($k < N$) such that z_k has critical sequence beginning $\zeta_k \mapsto \zeta_{k+1} \mapsto \zeta_{k+2} \mapsto \zeta_{k+3}$ for each $k < N$, where $\zeta_{N+2} = \eta_0$. Then there exist critical points ε_i ($i \leq C_{2N+2}(n)$) and embeddings e_i ($i < C_{2N+2}(n)$) such that e_i has critical sequence beginning $\zeta_{N+1} \mapsto \varepsilon_i \mapsto \varepsilon_{i+1}$ for each i , $\varepsilon_0 = \eta_0$, and $\varepsilon_{C_{2N+2}(n)} = \eta_n$.*

Proof. Proceed by induction on N ; we will prove the result for N , assuming it is true for $N' < N$. The induction step for N is similar to the proof of Lemma 4. Induct on n ; the case $n = 0$ is essentially vacuous (just let $\varepsilon_0 = \eta_0$). Given the result for n , if one has η_i for $i \leq n + 1$ and h_i for $i < n + 1$ as above, then the induction hypothesis gives critical points ε_i ($i \leq C_{2N+2}(n)$) and embeddings e_i ($i < C_{2N+2}(n)$) as required, with $\varepsilon_0 = \eta_0$ and $\varepsilon_{C_{2N+2}(n)} = \eta_n$. We can now apply Lemma 5, with $\zeta = \zeta_{N-1}$, $\zeta' = \zeta_N$, $\zeta'' = \zeta_{N+1}$, and $z = z_{N-1}$, to get critical points δ_i ($i \leq m$) and embeddings d_i ($i < m$), where $m = 2^{C_{2N+2}(n)}$, such that d_i has critical sequence beginning $\zeta_{N-1} \mapsto \zeta_N \mapsto \delta_i \mapsto \delta_{i+1}$ for each i , $\delta_0 = \zeta_{N+1}$, and $\delta_m = \varepsilon_{C_{2N+2}(n)} = \eta_n$. Finally, apply either Lemma 4 (if $N = 1$) or the induction hypothesis for $N - 1$ (if $N > 1$) to get critical points β_i ($i \leq C_{2N}(m)$) and embeddings b_i ($i < C_{2N}(m)$) such that b_i has critical sequence beginning $\zeta_N \mapsto \beta_i \mapsto \beta_{i+1}$ for each i , $\beta_0 = \delta_0 = \zeta_{N+1}$, and $\beta_{C_{2N}(m)} = \delta_m = \eta_n$. Define $\varepsilon_{C_{2N+2}(n)+i}$ for $1 \leq i \leq C_{2N}(m)$ by the equation $\varepsilon_{C_{2N+2}(n)+i} = h_n(\beta_i)$; note that this equation also holds for $i = 0$, since $h_n(\zeta_{N+1}) = \eta_n$. Hence, if we let $e_{C_{2N+2}(n)+i} = h_n(b_i)$ for $0 \leq i < C_{2N}(m)$, then $e_{C_{2N+2}(n)+i}$ has critical sequence beginning $\zeta_{N+1} \mapsto \varepsilon_{C_{2N+2}(n)+i} \mapsto \varepsilon_{C_{2N+2}(n)+i+1}$ for each such i . Also, $\varepsilon_{C_{2N+2}(n)+1} = h_n(\eta_n) = \eta_{n+1}$. This completes both inductions. ■

We are now ready to put everything together to get a lower bound on $F(n)$, the number of critical points below κ_n . Let us define $C_k(n)$ to be 2^n if n is not even and positive. In other words, the full definition of $C_k(n)$ is:

$$\begin{aligned} \text{If } k \text{ is 0 or odd:} & & C_k(n) &= 2^n \\ \text{If } k = 2: & & C_k(0) &= 0 \\ & & C_k(n+1) &= C_k(n) + 2^{C_k(n)} \\ \text{If } k \text{ is even and greater than 2:} & & C_k(0) &= 0 \\ & & C_k(n+1) &= C_k(n) + C_{k-2}(2^{C_k(n)}) \end{aligned}$$

Proposition 7. *Let $N \geq 0$, and suppose that one has critical points $\zeta_0 < \zeta_1 < \dots < \zeta_{N+1} < \eta_0 < \eta_1 < \dots < \eta_n$ and embeddings $h_i \in \mathcal{P}_j$ for $i < n$ such that h_i has critical sequence beginning $\zeta_N \mapsto \zeta_{N+1} \mapsto \eta_i \mapsto \eta_{i+1}$ for each i . Also suppose that there exist embeddings z_k ($k < N$) such that z_k has critical sequence beginning $\zeta_k \mapsto \zeta_{k+1} \mapsto \zeta_{k+2} \mapsto \zeta_{k+3}$ for each $k < N$, where $\zeta_{N+2} = \eta_0$. Then the number of critical points below η_n is at least $C_0(C_1(\dots(C_{2N+2}(n))\dots))$.*

Proof. Induct on N . For $N = 0$, the case $n = 0$ is trivial: $C_0(C_1(C_2(n))) = 2$ and $\zeta_0 < \zeta_1 < \eta_0$. If $N = 0$ and $n > 0$, apply Lemma 4 to get embeddings b_i and critical points β_i so that b_i has critical sequence beginning $\zeta_1 \mapsto \beta_i \mapsto \beta_{i+1}$ and $\beta_0 = \eta_0$, $\beta_{C_2(n)} = \eta_n$; then apply Lemma 3 (with $z = h_0$) to get embeddings a_i and critical points α_i so that a_i has critical sequence beginning $\zeta_0 \mapsto \alpha_i \mapsto \alpha_{i+1}$ and $\alpha_{C_1(C_2(n))} = \eta_n$; then apply Lemma 2 to get $C_0(C_1(C_2))$ critical points below η_n . For the induction step $N > 0$, given η_i and h_i as above, apply Lemma 6 to get embeddings e_i and critical points ε_i so that e_i has critical sequence beginning $\zeta_{N+1} \mapsto \varepsilon_i \mapsto \varepsilon_{i+1}$ and $\varepsilon_0 = \eta_0$, $\varepsilon_{C_{2N+2}(n)} = \eta_n$; then apply Lemma 5 (with $z = z_{N-1}$) to get embeddings d_i and critical points δ_i so that d_i has critical sequence beginning $\zeta_{N-1} \mapsto \zeta_N \mapsto \delta_i \mapsto \delta_{i+1}$ and $\delta_0 = \zeta_{N+1}$, $\delta_{C_{2N+1}(C_{2N+2}(n))} = \eta_n$; finally, apply the induction hypothesis for $N - 1$ to get $C_0(C_1(\dots(C_{2N+2}(n))\dots))$ critical points below η_n . ■

Now define the function g as follows: if $m \leq 2$, let $g(m) = m$; if $m \geq 3$, let $g(m) = C_0(C_1(\dots(C_{2m-4}(1))\dots))$.

Corollary 8. *Suppose one has critical points $\zeta_0 < \zeta_1 < \dots < \zeta_m$ and embeddings z_k ($0 \leq k \leq m - 3$) such that z_k has critical sequence beginning $\zeta_k \mapsto \zeta_{k+1} \mapsto \zeta_{k+2} \mapsto \zeta_{k+3}$. Then the number of critical points below ζ_m is at least $g(m)$.*

Proof. For $m < 3$, this is trivial. For $m \geq 3$, apply Proposition 7 with $N = m - 3$, $n = 1$, $\eta_0 = \zeta_{m-1}$, $\eta_1 = \zeta_m$, and $h_0 = z_{m-3}$. ■

Clearly we can apply Corollary 8 with $\zeta_k = \kappa_k$ and $z_k = j^{[k]}$ to get $F(m) \geq g(m)$ for all m . To complete the proof of Theorem 1, it will suffice to show that $g(m) \geq F_\omega(m - 1)$ for $m \geq 5$. (In fact, this is true for $m = 4$ as well: direct computation shows that $g(4) = 256$ and $F_\omega(3) = 61$.)

It is easy to prove by induction on N that $C_{2N}(n)$ is a strictly increasing function of n for each fixed N , and hence $C_{2N}(n) \geq n$. Of course, the same statements trivially hold for C_{2N+1} .

We can now prove by double induction on N and n that $C_{2N}(n+4) > F_{N+3}(n) + 3$ for all $N > 0$ and n . For $N = 1$ and $n = 0$, compute directly that $C_2(4) = 2059 > 16 = F_4(0) + 3$. If $N = 1$ and $n > 0$, then we have

$$C_2(n+4) = C_2(n+3) + 2^{C_2(n+3)} > 2^{C_2(n+3)} > 2^{F_4(n-1)+3} = F_4(n) + 3.$$

For $N > 1$ and $n = 0$, we have

$$C_{2N}(4) = C_{2N}(3) + C_{2N-2}(2^{C_{2N}(3)}) > C_{2N-2}(2^3) > C_{2N-2}(5) > F_{N+2}(1) + 3 = F_{N+3}(0) + 3.$$

Finally, if $N > 1$ and $n > 0$, then

$$\begin{aligned} C_{2N}(n+4) &= C_{2N}(n+3) + C_{2N-2}(2^{C_{2N}(n+3)}) > C_{2N-2}(C_{2N}(n+3) + 1) \\ &> C_{2N-2}(F_{N+3}(n-1) + 4) > F_{N+2}(F_{N+3}(n-1)) + 3 = F_{N+3}(n) + 3. \end{aligned}$$

Next, one can show by easy inductions on N that $C_{2N}(1) = 1$ and $C_{2N}(2) = N + 2$ for all $N \geq 1$. We are now ready to prove the desired bound on $g(m)$ for $m \geq 5$:

$$\begin{aligned} g(m) &= C_0(C_1(\dots(C_{2m-4}(1))\dots)) \\ &\geq C_{2m-8}(C_{2m-7}(C_{2m-6}(C_{2m-5}(C_{2m-4}(1))))) \\ &= C_{2m-8}(2^{m-1}) \\ &> C_{2m-8}(m+3) > F_{m-1}(m-1) = F_\omega(m-1). \end{aligned}$$

This completes the proof of Theorem 1.

The following remarks may help to clarify what is going on in the proof of Theorem 1. (Or they may make it more confusing. Read on at your own risk.)

First, let us observe a property which is not hard to see from the proofs of Lemmas 3–6 but which does not come out in the statements of these lemmas. Lemma 3 states that, given a sequence of embeddings b_i ($i < n$) such that b_i has critical sequence beginning $\zeta' \mapsto \beta_i \mapsto \beta_{i+1}$ (and an additional embedding z), one can obtain embeddings a_i ($i < 2^n$) such that a_i has critical sequence beginning $\zeta \mapsto \alpha_i \mapsto \alpha_{i+1}$, where $\alpha_{2^n} = \beta_n$. Now, if one can obtain some additional embeddings $b_n, b_{n+1}, \dots$ to extend the given sequence of b_i 's, then one gets a longer sequence of embeddings a_i , and it is clear from the proof of Lemma 3 that this longer sequence will extend the sequence of 2^n a_i 's obtained from $b_0, \dots, b_{n-1}$. Applying this repeatedly, we see that, if 2^m embeddings a_i are obtained from m embeddings b_i , then we will actually have $\alpha_{2^i} = \beta_i$ for all $i \leq m$, not just for $i = m$. Also, if we somehow obtained an infinite sequence of embeddings b_i , then we would get an infinite sequence of embeddings a_i and we would have $\alpha_{2^i} = \beta_i$ for all i . Similar statements hold for the situations in Lemmas 4–6.

Let us now introduce a more uniform notation. Instead of referring to embeddings a_i, b_i and critical points $\zeta, \zeta', \alpha_i, \beta_i$ in Lemma 3, one can instead state the result as follows: given embeddings $a_i^{(2)}$ ($i < n$) such that $a_i^{(2)}$ has critical sequence beginning $\zeta_1 \mapsto \alpha_i^{(2)} \mapsto \alpha_{i+1}^{(2)}$, and given that z_0 has critical sequence beginning $\zeta_0 \mapsto \zeta_1 \mapsto \alpha_0^{(2)}$, one can produce embeddings $a_i^{(1)}$ for $i < 2^n$ such that $a_i^{(1)}$ has critical sequence beginning $\zeta_1 \mapsto \alpha_i^{(1)} \mapsto \alpha_{i+1}^{(1)}$, where $\alpha_0^{(1)} = \zeta_1$ and $\alpha_{2^n}^{(1)} = \alpha_n^{(2)}$ (in fact, by the preceding paragraph, $\alpha_{2^i}^{(1)} = \alpha_i^{(2)}$ for all $i \leq n$). Similarly, one can write Lemma 4 so that it produces embeddings $a_i^{(2)}$ and critical points $\alpha_i^{(2)}$ from given embeddings $a_i^{(3)}$ and critical points $\alpha_i^{(3)}$ (writing the two fixed ordinals as ζ_0, ζ_1 instead of ζ, ζ'). Lemma 5 can be stated so that one uses $a_i^{(2N+2)}$ and $\alpha_i^{(2N+2)}$ to produce $a_i^{(2N+1)}$ and $\alpha_i^{(2N+1)}$, given fixed ordinals $\zeta_{N-1}, \zeta_N, \zeta_{N+1}$ and an initial embedding z_{N-1} . (One needs a variable superscript because Lemma 5 is applied many times, once for each inductive step in the proofs of Lemma 6 and Proposition 7.) Finally,

Lemma 6 produces embeddings $a_i^{(2N+2)}$ and critical points $\alpha_i^{(2N+2)}$ from given embeddings $a_i^{(2N+3)}$ and critical points $\alpha_i^{(2N+3)}$.

The reason for making these notational changes is so that one can think of Lemmas 3–6 as producing individual columns in a large two-dimensional array of embeddings, as shown in Figure 1. In this figure, each embedding is represented by giving the relevant part of its critical sequence. The boxed embeddings at the top of certain columns are the embeddings z_N which have to be given in advance; z_N appears at the top of column $2N + 3$, and z_0 also appears at the top of column 1. The unboxed embeddings in column N are obtained by applying the embeddings in column $N + 1$ to embeddings that have already been constructed; the lines drawn in the diagram connect each embedding in column $N + 1$ to the embeddings it is used to produce in column N .

One can now state Lemmas 3–6 in a unified form, as follows: If one is given the first n embeddings in column $N + 1$, as well as all of the boxed embeddings in preceding columns, then one can produce $C_N(n)$ embeddings in column N , ending up at the same critical point. Proposition 7 just combines these column-by-column results: given n embeddings in column $2N + 3$, one can produce $C_{2N+2}(n)$ embeddings in column $2N + 2$, $C_{2N+1}(C_{2N+2}(n))$ embeddings in column $2N + 1$, and so on, leading to $C_1(C_2(\dots(C_{2N+2}(n))\dots))$ embeddings in column 1 and hence, by Lemma 2, $C_0(C_1(\dots(C_{2N+2}(n))\dots))$ critical points.

The process of constructing new embeddings $a_i^{(N)}$ from old ones depends somewhat on the column number N . If N is odd, then the process is relatively simple and self-contained: given the first 2^n embeddings in column N , just apply $a_n^{(N+1)}$ to them to get the next 2^n embeddings. For $N = 2$, there is a little more work: given $C_2(n)$ embeddings in column 2, one must first run through the iteration for column 1 to produce $2^{C_2(n)}$ embeddings there, and then apply $a_n^{(3)}$ to the embeddings in column 1 to get $2^{C_2(n)}$ new embeddings in column 2. Finally, for column $2N$ where $N > 1$, one has to take two steps backward: given $C_{2N}(n)$ embeddings in column $2N$, one must produce the corresponding $2^{C_{2N}(n)}$ embeddings in column $2N - 1$ and $C_{2N-2}(2^{C_{2N}(n)})$ embeddings in column $2N - 2$ before applying $a_n^{(2N+1)}$ to get new embeddings in column $2N$. In this case, though, the process of computing embeddings in column $2N - 2$ from embeddings in column $2N - 1$ requires going back to columns $2N - 3$ and $2N - 4$, and so on; in fact, one must do computations all the way down to column 1 in order to produce the new embeddings for column $2N$.

One can think of the entire construction of Figure 1 as being a set of recursive equations for the embeddings $a_i^{(N)}$:

$$a_0^{(1)} = z_0; \quad a_0^{(2N+3)} = z_N;$$

if $C_N(n) \leq i < C_N(n + 1)$, then


$$a_i^{(N)} = a_n^{(N+1)}(a_{\bar{i}}^{(\hat{N})}),$$

where

$$\bar{i} = i - C_N(n),$$

$$\hat{N} = \begin{cases} N & \text{if } N \text{ is odd,} \\ N - 1 & \text{if } N = 2, \\ N - 2 & \text{if } N > 2 \text{ is even.} \end{cases}$$

This is a complicated recursion, where $a_i^{(N)}$ depends on embeddings in both earlier and later columns; hence, if one were to take this as the definition of the embeddings $a_i^{(N)}$, then one would have to check carefully that the recursion terminates after a finite number of steps to give a

FIGURE 1. The embeddings $a_i^{(N)}$ for $1 \leq N \leq 6$, $0 \leq i \leq 7$.

definition for $a_i^{(N)}$, and that the embeddings fit together in the proper way. This can be done, but it is rather messy, because the order in which the pairs (N, i) are processed is complicated. Perhaps the simplest description of this order is in computer science terms: the lines in Figure 1 define a collection of trees, one rooted at each boxed entry, and the recursion proceeds through a preorder traversal of the trees.

One final note: We have now shown that the number $F(n)$ of critical points below κ_n grows very rapidly, but this does not immediately imply that the number of critical points between κ_n and κ_{n+1} (which is $F(n+1) - F(n) - 1$) is large for *all* large n ; it is conceivable that F grows in spurts. (The construction in the next section makes this seem quite plausible.) This is not the case, however; the preceding construction shows that there are at least $g(n+1) - g(n) - 1$ critical points between ζ_n and ζ_{n+1} for $n \geq 2$. In fact, for the case $\zeta_n = \kappa_n$, if we let $h(n) = C_1(C_2(\dots(C_{2n-4}((1))\dots))$, then applying $a_{h(n)}^{(1)}$ to the $F(n)$ critical points below κ_n gives $F(n)$ additional critical points below $\alpha_{h(n)+1}^{(1)}$; we can now apply $a_{h(n)+1}^{(1)}$ to the current $2F(n)$ critical points to get $2F(n)$ more, and so on. (Here we are using the fact that all of the critical points below κ_n are in the interval $[\kappa_0, \kappa_n]$; this is true because, using the formula $cr(i_1(i_2)) = i_1(cr(i_2))$, one can show by an easy induction on k that $cr(k) \geq \kappa_0$ for all $k \in \mathcal{A}_j$ and hence for all $k \in \mathcal{P}_j$.) This shows that in fact $F(n+1) \geq 2^{h(n+1)-h(n)}F(n)$ for $n \geq 2$, so $F(n+1) - F(n)$ is much larger than $F(n)$.

3. USING VECTORS OF ORDINALS IN THE MAIN CONSTRUCTION

Theorem 1 and the growth rate estimate $F(n) > F_\omega(n-1)$ were obtained by applying Corollary 8 in the case where $\zeta_n = \kappa_n$. If we can find a sequence of critical points ζ_n growing more slowly than κ_n such that embeddings as described in Corollary 8 exist, then we get an improved lower bound on the growth rate of $F(n)$. In this section, we will obtain such sequences by another application of the same construction. We will again need the functions C_N and g from the preceding section.

The goal is to produce critical points ζ_i and embeddings z_i such that z_i has critical sequence beginning $\zeta_i \mapsto \zeta_{i+1} \mapsto \zeta_{i+2} \mapsto \zeta_{i+3}$. One can compare this with Lemma 3, in which one produces embeddings a_i and critical points α_i such that a_i has critical sequence beginning $\zeta \mapsto \alpha_i \mapsto \alpha_{i+1}$. The main difference here does not seem to be the fixed ordinal ζ ; after all, Lemma 5 differs from Lemma 3 only in having two fixed ordinals instead of one, and it is easy to prove a variant which has no fixed ordinal at all (in fact, this will look very much like Lemma 2). Rather, the main new difficulty is that successive embeddings must share three varying ordinals rather than just one. It turns out that one can get around this difficulty by considering ordinals in triples rather than individually.

More generally, we will consider finite sequences (‘vectors’) of ordinals, which we will always assume are nonempty and in strictly increasing order. If $\vec{\alpha}$ and $\vec{\beta}$ are two such sequences, $\vec{\alpha} < \vec{\beta}$ means that the last member of $\vec{\alpha}$ is less than the first member of $\vec{\beta}$. If $\vec{\alpha}$ is such a vector of length L , then its members will be denoted $\vec{\alpha}(0), \vec{\alpha}(1), \dots, \vec{\alpha}(L-1)$.

Given an embedding b , say that b has a critical vector sequence beginning $\vec{\beta}_0 \mapsto \vec{\beta}_1 \mapsto \dots \mapsto \vec{\beta}_n$ if $\vec{\beta}_0, \dots, \vec{\beta}_n$ are finite increasing sequences of ordinals of the same length, $\vec{\beta}_0 < \vec{\beta}_1 < \dots < \vec{\beta}_n$, the critical point of b is $\vec{\beta}_0(0)$, and $b(\vec{\beta}_k) = \vec{\beta}_{k+1}$ for each $k < n$. (One can think of “ $b(\vec{\beta}_k) = \vec{\beta}_{k+1}$ ” as an abbreviation for “ $b(\vec{\beta}_k(i)) = \vec{\beta}_{k+1}(i)$ for all i ” if one wants to apply embeddings only to ordinals and other embeddings.) Unlike ordinary critical sequences, a critical vector sequence is not uniquely defined; the ordinals $\vec{\beta}_0(i)$ for $i \geq 1$ can be arbitrary ordinals between $\vec{\beta}_0(0)$ and $\vec{\beta}_1(0)$. In the applications below, these additional ordinals will also be critical points.

Lemma 9. *Suppose that one has vectors $\vec{\tau} < \vec{\alpha}_0 < \vec{\alpha}_1 < \dots < \vec{\alpha}_n$ of length 3 and embeddings*

$a_i \in \mathcal{P}_j$ for $i < n$ such that a_i has a critical vector sequence beginning $\vec{\tau} \mapsto \vec{\alpha}_i \mapsto \vec{\alpha}_{i+1}$ for each i . Also suppose that there exist three embeddings z_0, z_1, z_2 which have critical sequences beginning $\vec{\tau}(0) \mapsto \vec{\tau}(1) \mapsto \vec{\tau}(2) \mapsto \vec{\alpha}_0(0)$, $\vec{\tau}(1) \mapsto \vec{\tau}(2) \mapsto \vec{\alpha}_0(0) \mapsto \vec{\alpha}_0(1)$, and $\vec{\tau}(2) \mapsto \vec{\alpha}_0(0) \mapsto \vec{\alpha}_0(1) \mapsto \vec{\alpha}_0(2)$, respectively. Then there exist critical points ζ_i ($i < 3(2^n + 1)$) and embeddings z_i ($i < 3 \cdot 2^n$) such that z_i has critical sequence beginning $\zeta_i \mapsto \zeta_{i+1} \mapsto \zeta_{i+2} \mapsto \zeta_{i+3}$ for each i , $\langle \zeta_0, \zeta_1, \zeta_2 \rangle = \vec{\tau}$, and $\langle \zeta_{3 \cdot 2^n}, \zeta_{3 \cdot 2^n + 1}, \zeta_{3 \cdot 2^n + 2} \rangle = \vec{\alpha}_n$.

Proof. Induct on n ; for $n = 0$, just let z_i be as given for $i = 0, 1, 2$, and let $\zeta_i = \vec{\tau}(i)$, $\zeta_{3+i} = \vec{\alpha}_0(i)$. Given the result for n , if one has $\vec{\alpha}_i$ for $i \leq n + 1$ and a_i for $i < n + 1$ as above, then the induction hypothesis gives ζ_i ($i < 3(2^n + 1)$) and z_i ($i < 3 \cdot 2^n$). Let $\zeta_{3 \cdot 2^n + i} = a_n(\zeta_i)$ for $3 \leq i < 3 \cdot 2^n + 3$; this equation holds also for $i = 0, 1, 2$, since $a_n(\vec{\tau}) = \vec{\alpha}_n$. Hence, if we let $z_{3 \cdot 2^n + i} = a_n(z_i)$ for $i < 3 \cdot 2^n$, then the embeddings $z_{3 \cdot 2^n + i}$ will have the required critical sequences. Finally, $\langle \zeta_{3 \cdot 2^n + 1}, \zeta_{3 \cdot 2^n + 2}, \zeta_{3 \cdot 2^n + 3} \rangle = a_n(\vec{\alpha}_n) = \vec{\alpha}_{n+1}$. This completes the induction. ■

We now have reduced the problem of obtaining embeddings z_i related as above to that of obtaining embeddings a_i which map triples of ordinals as above. But it is not hard to see that the methods used for Lemmas 3–6 work without change if one uses vectors of ordinals (of a fixed finite length) instead of individual ordinals. For instance, the ‘vectorized’ version of Lemma 3 is:

Lemma 10. *Suppose that one has vectors $\vec{\tau} < \vec{\tau}' < \vec{\beta}_0 < \vec{\beta}_1 < \dots < \vec{\beta}_n$ and embeddings $b_i \in \mathcal{P}_j$ for $i < n$ such that b_i has a critical vector sequence beginning $\vec{\tau}' \mapsto \vec{\beta}_i \mapsto \vec{\beta}_{i+1}$ for each i . Also suppose that there is an embedding t such that t has a critical vector sequence beginning $\vec{\tau} \mapsto \vec{\tau}' \mapsto \vec{\beta}_0$. Then there exist vectors $\vec{\alpha}_i$ ($i \leq 2^n$) and embeddings a_i ($i < 2^n$) such that a_i has a critical vector sequence beginning $\vec{\tau} \mapsto \vec{\alpha}_i \mapsto \vec{\alpha}_{i+1}$ for each i , $\vec{\alpha}_0 = \vec{\tau}'$, and $\vec{\alpha}_{2^n} = \vec{\beta}_n$.*

The proof of this lemma is the same as that of the original Lemma 3. (Here we have $b_n(\vec{\tau}) = \vec{\tau}$ because all of the ordinals $\vec{\tau}(i)$ are below $\vec{\tau}'(0)$, which is the critical point of b_n .) It is just as easy to produce vector versions of Lemmas 4–6. One can now combine these as before to get the following analogue of Corollary 8:

Lemma 11. *Suppose one has vectors $\vec{\tau}_0 < \vec{\tau}_1 < \dots < \vec{\tau}_m$ of length 3 and embeddings t_k ($0 \leq k \leq m - 3$) such that t_k has a critical vector sequence beginning $\vec{\tau}_k \mapsto \vec{\tau}_{k+1} \mapsto \vec{\tau}_{k+2} \mapsto \vec{\tau}_{k+3}$. Suppose also that there exist three embeddings z_0, z_1, z_2 which have critical sequences beginning $\vec{\tau}_0(0) \mapsto \vec{\tau}_0(1) \mapsto \vec{\tau}_0(2) \mapsto \vec{\tau}_1(0)$, $\vec{\tau}_0(1) \mapsto \vec{\tau}_0(2) \mapsto \vec{\tau}_1(0) \mapsto \vec{\tau}_1(1)$, and $\vec{\tau}_0(2) \mapsto \vec{\tau}_1(0) \mapsto \vec{\tau}_1(1) \mapsto \vec{\tau}_1(2)$, respectively. Then there exist critical points ζ_i ($i < 3(g(n) + 1)$) and embeddings z_i ($i < 3g(n)$) such that z_i has critical sequence beginning $\zeta_i \mapsto \zeta_{i+1} \mapsto \zeta_{i+2} \mapsto \zeta_{i+3}$ for each i , $\langle \zeta_0, \zeta_1, \zeta_2 \rangle = \vec{\tau}_0$, and $\langle \zeta_{3g(m)}, \zeta_{3g(m)+1}, \zeta_{3g(m)+2} \rangle = \vec{\tau}_m$.*

It follows that, if the hypotheses of Lemma 11 hold, then the number of critical points below $\vec{\tau}_m(0)$ is at least $g(3g(m))$. For instance, if we let $\vec{\tau}_k = \langle \kappa_{3k}, \kappa_{3k+1}, \kappa_{3k+2} \rangle$, $t_k = j^{[3k]} \circ j^{[3k]} \circ j^{[3k]}$, and $z_i = j^{[i]}$ for $i = 0, 1, 2$, then we get $F(3m) \geq g(3g(m))$, which is a much better bound than $g(3m)$ for $m \geq 3$.

The hypotheses of Lemma 11 look very similar to those of Corollary 8; one merely has to have suitable embeddings working on triples of ordinals rather than ordinals. Hence, in order to meet these hypotheses, we merely have to repeat the modifications above, replacing triples of ordinals with triples of triples of ordinals, or, equivalently, vectors of length 9. More generally, the proof of Lemma 11 goes through without change to produce the following version:

Lemma 12. *Suppose one has vectors $\vec{\tau}_0 < \vec{\tau}_1 < \dots < \vec{\tau}_m$ of length $3L$ and embeddings t_k ($0 \leq k \leq m - 3$) such that t_k has a critical vector sequence beginning $\vec{\tau}_k \mapsto \vec{\tau}_{k+1} \mapsto \vec{\tau}_{k+2} \mapsto \vec{\tau}_{k+3}$. Let*

$\vec{\tau}_k^{[i]}$ for $i = 0, 1, 2$ be the three sequences of length L composing $\vec{\tau}_k$ (i.e., $\vec{\tau}_k = \vec{\tau}_k^{[0]} \cap \vec{\tau}_k^{[1]} \cap \vec{\tau}_k^{[2]}$), and suppose that there exist three embeddings z_0, z_1, z_2 which have critical vector sequences beginning $\vec{\tau}_0^{[0]} \mapsto \vec{\tau}_0^{[1]} \mapsto \vec{\tau}_0^{[2]} \mapsto \vec{\tau}_1^{[0]}, \vec{\tau}_0^{[1]} \mapsto \vec{\tau}_0^{[2]} \mapsto \vec{\tau}_1^{[0]} \mapsto \vec{\tau}_1^{[1]},$ and $\vec{\tau}_0^{[2]} \mapsto \vec{\tau}_1^{[0]} \mapsto \vec{\tau}_1^{[1]} \mapsto \vec{\tau}_1^{[2]}$, respectively. Then there exist vectors $\vec{\zeta}_i$ ($i < 3(g(n) + 1)$) and embeddings z_i ($i < 3g(n)$) such that z_i has a critical vector sequence beginning $\vec{\zeta}_i \mapsto \vec{\zeta}_{i+1} \mapsto \vec{\zeta}_{i+2} \mapsto \vec{\zeta}_{i+3}$ for each i , $\vec{\zeta}_0 \cap \vec{\zeta}_1 \cap \vec{\zeta}_2 = \vec{\tau}_0$, and $\vec{\zeta}_{3g(m)} \cap \vec{\zeta}_{3g(m)+1} \cap \vec{\zeta}_{3g(m)+2} = \vec{\tau}_m$.

Actually, we have $\vec{\zeta}_{3g(k)} \cap \vec{\zeta}_{3g(k)+1} \cap \vec{\zeta}_{3g(k)+2} = \vec{\tau}_k$ for all $k \leq m$.

By using Lemma 12 repeatedly, we can generate critical points according to iterates of the function g . Let $g_m(n) = g((3g)^m(n))$; that is, $g_0(n) = g(n)$ and $g_{m+1}(n) = g_m(3g(n))$. (One also has $g_{m+1}(n) = g(3g_m(n))$.) For any fixed m , a suitable starting point for applying Lemma 12 is the sequences $\vec{\tau}_n = \langle \kappa_{3^m n+i} : 0 \leq i < 3^m \rangle$, since we can define t_n to be $j^{3^m n}(j^{3^m})$, where j^l is the composition of l j 's. The three additional embeddings z_i are just $j^{3^{m-1}i}(j^{3^{m-1}})$, $i = 0, 1, 2$. Lemma 12 now lets us define z_n and $\vec{\zeta}_n$ (of length 3^{m-1}) for all n , and we have $\vec{\zeta}_{3g(n)}(0) = \vec{\tau}_n(0) = \kappa_{3^m n}$. Furthermore, we have $\vec{\zeta}_n = \langle \kappa_{3^{m-1}n+i} : 0 \leq i < 3^{m-1} \rangle$ for $n \leq 5$ (for $n \leq 8$, actually, but this is irrelevant), so we can define three embeddings $z'_i = j^{3^{m-2}i}(j^{3^{m-2}})$, $i = 0, 1, 2$, and again we are in a position to apply Lemma 12. Continuing this through a total of $m - 1$ applications of Lemma 12 leaves us with ordinal sequences of length 3, and we are now in a position to apply Lemma 11 and then Corollary 8. We therefore get the following result: for all m and n , $F(3^m n) \geq g_m(n)$.

In particular, we have $F(3^m) \geq g_m(1)$ for all m . Now, a direct computation shows that $g_1(1) = 4 > 3 = F_{\omega+1}(0)$. This, together with the equation $g_{m+1}(n) = g(3g_m(n))$ and the inequality $g(n) > F_{\omega+1}(n - 1)$ for $n \geq 4$ from the preceding section, yields an easy inductive proof that $g_m(1) > F_{\omega+1}(m - 1)$ for all $m \geq 1$. So $F(3^m) > F_{\omega+1}(m - 1)$. If we apply this in the case where $m = \lfloor \log_3(n) \rfloor$, we get the result stated in the introduction: $F(n) > F_{\omega+1}(\lfloor \log_3(n) \rfloor - 1)$ for all $n \geq 3$.

4. IRREGULAR CONSTRUCTIONS

The preceding sections have shown that $F(n)$, the number of critical points below κ_n , is very large when n is large. Now we will turn to the specific value $F(4)$. (This is the first value of F that could be large; we will note in the next section that $F(3) = 4$.) The lower bound for $F(4)$ given by section 2 is $g(4) = 256$, and section 3 does not improve this value. Here we will show, by use of some *ad hoc* modifications of the main construction, that $F(4)$ is quite large.

The numerical functions involved will be somewhat different from those used in section 2; the new functions will be called $\tilde{C}_N$ rather than C_N . Also, the resulting critical points and embeddings will be referred to as $\tilde{\alpha}_i^{(N)}$ and $\tilde{a}_i^{(N)}$ rather than $\alpha_i^{(N)}$ and $a_i^{(N)}$ as at the end of section 2.

The first four stages of the construction, Lemmas 2–5 (or the first four columns of Figure 1), will remain unchanged: $\tilde{C}_N$ is the same as C_N for $N \leq 3$. Since this is a specific construction, we will go ahead and use the specific values $\kappa_0, \kappa_1, \kappa_2, \kappa_3$ rather than the general $\zeta_0, \zeta_1, \zeta_2, \zeta_3$. In other words, we will use Lemma 5 with $\zeta = \kappa_0$, $\zeta' = \kappa_1$, $\zeta'' = \kappa_2$, and $z = j$.

As a first example, suppose that we have an embedding with critical point κ_2 which sends κ_2 to κ_3 and κ_3 to some critical point μ . Then we can apply Lemma 5 with $n = 1$, followed by Lemmas 4, 3, and 2, to produce $2^{2^{C_2(2^1)}} = 256$ critical points below μ . Hence, if μ is less than κ_4 , then $F(4) > 256$.

We will see in the next section that such an embedding exists, with $\mu < \kappa_4$; in fact, $j_{(12)}$ will work, where $j_{(n)}$ is as defined in section 1. For the rest of this section, we will continue to construct

many critical points below κ_4 , assuming that we can get certain specific embeddings; the proofs that such embeddings exist will be given in the next section.

For the next step, we will construct two additional stages (columns) by a method different from that used in section 2. First, we give the recursive definitions of the relevant functions $\tilde{C}_4$ and $\tilde{C}_5$:

$$\begin{aligned}\tilde{C}_4(0) &= 0, & \tilde{C}_4(1) &= 8, & \tilde{C}_4(m+1) &= \tilde{C}_4(m) + \tilde{C}_1(\tilde{C}_2(\tilde{C}_3(\tilde{C}_4(m)))) - 1 \quad \text{for } m > 0; \\ \tilde{C}_5(0) &= 0, & \tilde{C}_5(1) &= 2, & \tilde{C}_5(m+1) &= \tilde{C}_5(m) + \tilde{C}_3(\tilde{C}_4(\tilde{C}_5(m))) \quad \text{for } m > 0.\end{aligned}$$

At this point, it will be useful to introduce an abbreviation: for $N < M$, let

$$\tilde{C}_{[N,M]}(n) = \tilde{C}_N(\tilde{C}_{N+1}(\dots(\tilde{C}_{M-1}(n))\dots)).$$

So the recursive formula for $\tilde{C}_4$ can be written as $\tilde{C}_4(m+1) = \tilde{C}_4(m) + \tilde{C}_{[1,4]}(\tilde{C}_4(m)) - 1$. (One could further abbreviate this to $\tilde{C}_4(m) + \tilde{C}_{[1,5]}(m) - 1$, but it seems preferable to keep the function being defined explicit.) And now the corresponding lemmas:

Lemma 13. *Suppose that one has critical points $\tilde{\alpha}_0^{(5)} < \tilde{\alpha}_1^{(5)} < \dots < \tilde{\alpha}_n^{(5)}$, where $n > 0$, and embeddings $\tilde{a}_i^{(5)}$ for $i < n$ such that $\tilde{a}_i^{(5)}$ has critical sequence beginning $\kappa_0 \mapsto \kappa_2 \mapsto \tilde{\alpha}_i^{(5)} \mapsto \tilde{\alpha}_{i+1}^{(5)}$ for each i . Also suppose that $\tilde{a}_0^{(5)}(\kappa_1) = \kappa_3$. Then there exist critical points $\tilde{\alpha}_i^{(4)}$ ($i \leq \tilde{C}_4(n)$) and embeddings $\tilde{a}_i^{(4)}$ ($i < \tilde{C}_4(n)$) such that $\tilde{a}_i^{(4)}$ has critical sequence beginning $\kappa_2 \mapsto \tilde{\alpha}_i^{(4)} \mapsto \tilde{\alpha}_{i+1}^{(4)}$ for each i , $\tilde{\alpha}_0^{(4)} = \kappa_3$, $\tilde{\alpha}_1^{(4)} = \tilde{\alpha}_0^{(5)}$, and $\tilde{\alpha}_{\tilde{C}_4(n)}^{(4)} = \tilde{\alpha}_n^{(5)}$.*

Proof. Induct on $n \geq 1$. For $n = 1$, first let $\tilde{a}_0^{(4)} = \tilde{a}_0^{(5)}(j)$, $\tilde{\alpha}_0^{(4)} = \kappa_3$, and $\tilde{\alpha}_1^{(4)} = \tilde{\alpha}_0^{(5)}$. Now apply Lemma 5 (with $z = j$), Lemma 4, and Lemma 3 (with $z = j$ again) to get $\tilde{C}_1(\tilde{C}_2(\tilde{C}_3(1))) = 8$ embeddings $\tilde{a}_i^{(1)}$ and corresponding critical points $\tilde{\alpha}_i^{(1)}$ ($i \leq 8$) such that $\tilde{a}_i^{(1)}$ has critical sequence beginning $\kappa_0 \mapsto \tilde{\alpha}_i^{(1)} \mapsto \tilde{\alpha}_{i+1}^{(1)}$, and $\tilde{\alpha}_8^{(1)} = \tilde{\alpha}_0^{(5)}$; since we used $z = j$ in Lemma 3, we will have $\tilde{a}_0^{(1)} = j$, $\tilde{\alpha}_0^{(1)} = \kappa_1$, and $\tilde{\alpha}_1^{(1)} = \kappa_2$. Now let $\tilde{a}_i^{(4)} = \tilde{a}_0^{(5)}(\tilde{a}_i^{(1)})$ and $\tilde{\alpha}_{i+1}^{(4)} = \tilde{a}_0^{(5)}(\tilde{\alpha}_{i+1}^{(1)})$ for $1 \leq i < 8$; these equations also hold for $i = 0$, and we get $\tilde{\alpha}_8^{(4)} = \tilde{\alpha}_1^{(5)}$, so everything fits together and the case $n = 1$ is done.

The induction step is similar. Given the result for n , if one has $\tilde{\alpha}_i^{(5)}$ for $i \leq n+1$ and $\tilde{a}_i^{(5)}$ for $i < n+1$ as above, then the induction hypothesis gives critical points $\tilde{\alpha}_i^{(4)}$ ($i \leq \tilde{C}_4(n)$) and embeddings $\tilde{a}_i^{(4)}$ ($i < \tilde{C}_4(n)$) as required, with $\tilde{\alpha}_0^{(4)} = \kappa_3$, $\tilde{\alpha}_1^{(4)} = \tilde{\alpha}_0^{(5)}$, and $\tilde{\alpha}_{\tilde{C}_4(n)}^{(4)} = \tilde{\alpha}_n^{(5)}$. We can now apply Lemmas 5, 4, and 3 successively to get embeddings $\tilde{a}_i^{(1)}$ ($i < \tilde{C}_{[1,4]}(\tilde{C}_4(n))$) and critical points $\tilde{\alpha}_i^{(1)}$ ($i \leq \tilde{C}_{[1,4]}(\tilde{C}_4(n))$) such that $\tilde{a}_i^{(1)}$ has critical sequence beginning $\kappa_0 \mapsto \tilde{\alpha}_i^{(1)} \mapsto \tilde{\alpha}_{i+1}^{(1)}$; also, we will have $\tilde{\alpha}_1^{(1)} = \kappa_2$ and $\tilde{\alpha}_{\tilde{C}_{[1,4]}(\tilde{C}_4(n))}^{(1)} = \tilde{\alpha}_n^{(5)}$. Now let $\tilde{a}_{\tilde{C}_4(n)-1+i}^{(4)} = \tilde{a}_n^{(5)}(\tilde{a}_i^{(1)})$ and $\tilde{\alpha}_{\tilde{C}_4(n)+i}^{(4)} = \tilde{a}_n^{(5)}(\tilde{\alpha}_{i+1}^{(1)})$ for $1 \leq i < \tilde{C}_{[1,4]}(\tilde{C}_4(n))$; this fits together suitably because $\tilde{a}_n^{(5)}(\tilde{\alpha}_1^{(1)}) = \tilde{\alpha}_n^{(5)} = \tilde{\alpha}_{\tilde{C}_4(n)}^{(4)}$, and we also get $\tilde{\alpha}_{\tilde{C}_4(n)+1}^{(4)} = \tilde{\alpha}_{n+1}^{(5)}$, so the induction is complete. $\blacksquare$

Lemma 14. *Suppose that one has critical points $\tilde{\alpha}_0^{(6)} < \tilde{\alpha}_1^{(6)} < \dots < \tilde{\alpha}_n^{(6)}$, where $n > 0$, and embeddings $\tilde{a}_i^{(6)}$ for $i < n$ such that $\tilde{a}_i^{(6)}$ has critical sequence beginning $\kappa_1 \mapsto \kappa_2 \mapsto \tilde{\alpha}_i^{(6)} \mapsto \tilde{\alpha}_{i+1}^{(6)}$ for each i . Also suppose that $\tilde{a}_0^{(6)}(j)(\kappa_1) = \kappa_3$. Then there exist critical points $\tilde{\alpha}_i^{(5)}$ ($i \leq \tilde{C}_5(n)$) and*

embeddings $\tilde{a}_i^{(5)}$ ($i < \tilde{C}_5(n)$) such that $\tilde{a}_i^{(5)}$ has critical sequence beginning $\kappa_0 \mapsto \kappa_2 \mapsto \tilde{\alpha}_i^{(5)} \mapsto \tilde{\alpha}_{i+1}^{(5)}$ for each i , $\tilde{a}_0^{(5)}(\kappa_1) = \kappa_3$, $\tilde{\alpha}_0^{(5)} = \tilde{\alpha}_0^{(6)}$, and $\tilde{\alpha}_{\tilde{C}_5(n)}^{(5)} = \tilde{\alpha}_n^{(6)}$.

Proof. Induct on n . For $n = 1$, let $\tilde{a}_0^{(5)} = \tilde{a}_0^{(6)}(j)$, $\tilde{a}_1^{(5)} = \tilde{a}_0^{(6)}(\tilde{a}_0^{(5)}(j)(j))$, $\tilde{\alpha}_0^{(5)} = \tilde{\alpha}_0^{(6)}$, $\tilde{\alpha}_1^{(5)} = \tilde{a}_0^{(6)}(\kappa_3)$, and $\tilde{\alpha}_2^{(5)} = \tilde{\alpha}_1^{(6)}$. [This is just a shorter way of stating the following: Let $\tilde{a}_0^{(5)} = \tilde{a}_0^{(6)}(j)$, $\tilde{\alpha}_0^{(5)} = \tilde{\alpha}_0^{(6)}$, and $\tilde{\alpha}_1^{(5)} = \tilde{a}_0^{(6)}(\kappa_3)$. Apply Lemma 13 with $n = 1$, and then apply Lemma 5 with $n = 1$ (not $n = 8$), to get embeddings $\tilde{a}_i^{(3)}$ ($i < 2$) and critical points $\tilde{\alpha}_i^{(3)}$ ($i \leq 2$) such that $\tilde{a}_i^{(3)}$ has critical sequence beginning $\kappa_0 \mapsto \kappa_1 \mapsto \tilde{\alpha}_i^{(3)} \mapsto \tilde{\alpha}_{i+1}^{(3)}$, and $\tilde{\alpha}_0^{(3)} = \kappa_2$, $\tilde{\alpha}_1^{(3)} = \kappa_3$, and $\tilde{\alpha}_2^{(3)} = \tilde{\alpha}_0^{(6)}$. Now let $\tilde{a}_1^{(5)} = \tilde{a}_0^{(6)}(\tilde{a}_1^{(3)})$ and $\tilde{\alpha}_2^{(5)} = \tilde{\alpha}_1^{(6)}$.]

Given the result for n , if one has $\tilde{\alpha}_i^{(6)}$ for $i \leq n + 1$ and $\tilde{a}_i^{(6)}$ for $i < n + 1$ as above, then the induction hypothesis gives critical points $\tilde{\alpha}_i^{(5)}$ ($i \leq \tilde{C}_5(n)$) and embeddings $\tilde{a}_i^{(5)}$ ($i < \tilde{C}_5(n)$) as required, with $\tilde{a}_0^{(5)}(\kappa_1) = \kappa_3$, $\tilde{\alpha}_0^{(5)} = \tilde{\alpha}_0^{(6)}$, and $\tilde{\alpha}_{\tilde{C}_5(n)}^{(5)} = \tilde{\alpha}_n^{(6)}$. We can now apply Lemmas 13 and 5 successively to get embeddings $\tilde{a}_i^{(3)}$ ($i < \tilde{C}_{[3,5]}(\tilde{C}_5(n))$) and critical points $\tilde{\alpha}_i^{(3)}$ ($i \leq \tilde{C}_{[3,5]}(\tilde{C}_5(n))$) such that $\tilde{a}_i^{(3)}$ has critical sequence beginning $\kappa_0 \mapsto \kappa_1 \mapsto \tilde{\alpha}_i^{(3)} \mapsto \tilde{\alpha}_{i+1}^{(3)}$; also, also, we will have $\tilde{\alpha}_0^{(3)} = \kappa_2$ and $\tilde{\alpha}_{\tilde{C}_{[3,5]}(\tilde{C}_5(n))}^{(3)} = \tilde{\alpha}_n^{(6)}$. Now let $\tilde{a}_{\tilde{C}_5(n)+i}^{(5)} = \tilde{a}_n^{(6)}(\tilde{a}_i^{(3)})$ and $\tilde{\alpha}_{\tilde{C}_5(n)+i+1}^{(5)} = \tilde{a}_n^{(6)}(\tilde{\alpha}_{i+1}^{(3)})$ for $0 \leq i < \tilde{C}_{[3,5]}(\tilde{C}_5(n))$; this fits together suitably because $\tilde{a}_n^{(6)}(\tilde{\alpha}_0^{(3)}) = \tilde{\alpha}_n^{(6)} = \tilde{\alpha}_{\tilde{C}_5(n)}^{(5)}$, and we also get $\tilde{\alpha}_{\tilde{C}_5(n)+1}^{(5)} = \tilde{\alpha}_{n+1}^{(6)}$, so the induction is complete. $\blacksquare$


The modified construction to this point is shown in Figure 2. (For now, ignore the references to $\tilde{\alpha}_i^{(7)}$.) The main difference between this and the original construction is that here we extend column 4 by applying embeddings in column 5 to those in column 1 instead of column 2, and we extend column 5 by applying embeddings in column 6 to those in column 3 instead of column 5.

We now assume the existence of an embedding k such that k has critical sequence beginning $\kappa_1 \mapsto \kappa_2 \mapsto \mu \mapsto \nu$ for some μ and ν , satisfying the additional condition $k(j)(\kappa_1) = \kappa_3$. (We will see in the next section that $k = j_{(10)}$ works.) Then we can let $\tilde{a}_0^{(6)} = k$, $\tilde{\alpha}_0^{(6)} = \mu$, and $\tilde{\alpha}_1^{(6)} = \nu$, and apply Lemmas 14, 13, 5, 4, 3, and 2 successively to fill in all of Figure 2 except the part below the dashed line; this generates a total of $\tilde{C}_{[0,6]}(1)$ critical points below ν .

Since we have $\tilde{C}_2 = C_2$, we can use the estimate $\tilde{C}_2(m+4) \geq F_4(m) + 3$. This leads to the computations $\tilde{C}_4(1) = 8$, $\tilde{C}_{[3,5]}(1) = 256$, $\tilde{C}_{[2,5]}(1) \geq F_4(252) + 3$, $\tilde{C}_{[1,5]}(1) > F_4(253)$, $\tilde{C}_4(2) > F_4(253) + 7$, $\tilde{C}_{[3,5]}(2) > F_4(254) + 4$, and $\tilde{C}_{[0,6]}(1) = \tilde{C}_{[0,5]}(2) > \tilde{C}_{[2,5]}(2) > F_4(F_4(254))$, so there are more than $F_4(F_4(254))$ critical points below ν . The value of ν produced in the next section will be less than κ_4 , so we will have $F(4) > F_4(F_4(254)) > F_5(1)$. This already shows that $F(4)$ is quite large.

To get even larger lower bounds on $F(4)$, we will add more columns to the irregular construction. To do this, we assume the existence of two additional embeddings k', k'' such that k' has critical sequence beginning $\kappa_0 \mapsto \mu \mapsto \nu \mapsto \xi$ (where μ and ν are the same as for the embedding k), k'' has critical sequence beginning $\kappa_2 \mapsto \nu$, and $k''(k)(j)(\mu) = \xi$. (In the next section, we will verify that the embeddings $k' = j_{(10)}(j_{(11)})$ and $k'' = j_{(9)}(j_{(14)})$ satisfy these conditions.)


The new part of the construction is shown in Figure 3. The boxed entries are the given embeddings k' and k'' . The new functions $\tilde{C}_N$ indicating how many embeddings in column N are

FIGURE 2. The embeddings $\tilde{\alpha}_m^{(n)}$ for $1 \leq n \leq 6$, $0 \leq m \leq 5$.

obtained from a given number of embeddings in column $N + 1$ are defined as follows:

$$\begin{aligned}
 \tilde{C}_6(m) &= 2^m; \\
 \tilde{C}_7(0) &= 0, \quad \tilde{C}_7(m+1) = \tilde{C}_7(m) + \tilde{C}_{[1,7]}(\tilde{C}_7(m)) - 8; \\
 \tilde{C}_8(m) &= 2^m; \\
 \tilde{C}_9(0) &= 0, \quad \tilde{C}_9(1) = \tilde{C}_{[1,9]}(\tilde{C}_{[1,6]}(1) - 8) - 8, \\
 \tilde{C}_9(m+1) &= \tilde{C}_9(m) + \tilde{C}_{[1,9]}(\tilde{C}_9(m)) - \tilde{C}_{[1,6]}(1) \quad \text{for } m > 0; \\
 \tilde{C}_{10}(0) &= 0, \quad \tilde{C}_{10}(m+1) = \tilde{C}_{10}(m) + \tilde{C}_{[3,10]}(\tilde{C}_{10}(m)).
 \end{aligned}$$

Instead of giving six more lemmas for the construction of columns 6–11, we will just give a description of how column number N is constructed from column $N + 1$ for $N = 6, 7, 8, 9, 10, 11$.

FIGURE 3. The embeddings $\tilde{a}_m^{(n)}$ for $7 \leq n \leq 12$, $0 \leq m \leq 5$.

From this, it is straightforward to state corresponding lemmas, and the proofs will be similar to those of preceding lemmas.

Column 6 is constructed from column 7 using Lemma 5; no new results are needed.

If one has already used n embeddings in column 8 to produce $\tilde{C}_7(n)$ embeddings in column 7 ending up with $\tilde{\alpha}_{\tilde{C}_7(n)}^{(7)} = \tilde{\alpha}_n^{(8)}$, then one can apply the preceding results to get $\tilde{C}_{[1,7]}(\tilde{C}_7(n))$ embeddings in column 1, and one will have $\tilde{\alpha}_8^{(1)} = \mu$ and $\tilde{\alpha}_{\tilde{C}_{[1,7]}(\tilde{C}_7(n))}^{(1)} = \tilde{\alpha}_n^{(8)}$. Now apply $\tilde{a}_n^{(8)}$ to all but the first 8 of these embeddings in column 1 to get $\tilde{C}_{[1,7]}(\tilde{C}_7(n)) - 8$ new embeddings for column 7, for a total of $\tilde{C}_7(n+1)$ embeddings ending up with $\tilde{\alpha}_{\tilde{C}_7(n+1)}^{(7)} = \tilde{\alpha}_{n+1}^{(8)}$.

Column 8 is constructed from column 9 using Lemma 5.

To build column 10 from column 9 requires an extra assumption not visible in Figure 3: $\tilde{a}_0^{(10)}(\mu) = \xi$. Given this, one can proceed as follows. First, use the given embedding k in col-

n	Critical sequence of $\tilde{a}_m^{(n)}$	$\tilde{a}_0^{(n)}$	$\tilde{\alpha}_0^{(n)}$	$\hat{n}$	$\tilde{C}_n(0), (1)$	$\tilde{C}_n(m+1) - \tilde{C}_n(m)$	Growth
0	$\tilde{\alpha}_m^{(0)} \mapsto \tilde{\alpha}_{m+1}^{(0)}$	j	κ_0	0	1	$\tilde{C}_0(m)$	$F_3(m-3)$
1	$\kappa_0 \mapsto \tilde{\alpha}_m^{(1)} \mapsto \tilde{\alpha}_{m+1}^{(1)}$	j	κ_1	1	1	$\tilde{C}_1(m)$	$F_3(m-3)$
2	$\kappa_1 \mapsto \tilde{\alpha}_m^{(2)} \mapsto \tilde{\alpha}_{m+1}^{(2)}$		κ_2	1	0	$\tilde{C}_1(\tilde{C}_2(m))$	$F_4(m-4)$
3	$\kappa_0 \mapsto \kappa_1 \mapsto \tilde{\alpha}_m^{(3)} \mapsto \tilde{\alpha}_{m+1}^{(3)}$	j	κ_2	3	1	$\tilde{C}_3(m)$	$F_3(m-3)$
4	$\kappa_2 \mapsto \tilde{\alpha}_m^{(4)} \mapsto \tilde{\alpha}_{m+1}^{(4)}$		κ_3	1	0, 8	$\tilde{C}_{[1,4]}(\tilde{C}_4(m)) - 1$	$F_5(m-2)$
5	$\kappa_0 \mapsto \kappa_2 \mapsto \tilde{\alpha}_m^{(5)} \mapsto \tilde{\alpha}_{m+1}^{(5)}$		μ	3	0, 2	$\tilde{C}_{[3,5]}(\tilde{C}_5(m))$	$F_6(m-3)$
6	$\kappa_1 \mapsto \kappa_2 \mapsto \tilde{\alpha}_m^{(6)} \mapsto \tilde{\alpha}_{m+1}^{(6)}$	k	μ	6	1	$\tilde{C}_6(m)$	$F_3(m-3)$
7	$\mu \mapsto \tilde{\alpha}_m^{(7)} \mapsto \tilde{\alpha}_{m+1}^{(7)}$		ν	1	0	$\tilde{C}_{[1,7]}(\tilde{C}_7(m)) - 8$	$F_7(m-2)$
8	$\kappa_0 \mapsto \mu \mapsto \tilde{\alpha}_m^{(8)} \mapsto \tilde{\alpha}_{m+1}^{(8)}$	k'	ν	8	1	$\tilde{C}_8(m)$	$F_3(m-3)$
9	$\nu \mapsto \tilde{\alpha}_m^{(9)} \mapsto \tilde{\alpha}_{m+1}^{(9)}$		ξ	1	0, [*]	$\tilde{C}_{[1,9]}(\tilde{C}_9(m)) - \tilde{C}_{[1,6]}(1)$	$F_8(m-2)$
10	$\kappa_0 \mapsto \nu \mapsto \tilde{\alpha}_m^{(10)} \mapsto \tilde{\alpha}_{m+1}^{(10)}$		$k''(k)(\kappa_2)$	3	0	$\tilde{C}_{[3,10]}(\tilde{C}_{10}(m))$	$F_9(m-2)$
11	$\kappa_1 \mapsto \nu \mapsto \tilde{\alpha}_m^{(11)} \mapsto \tilde{\alpha}_{m+1}^{(11)}$		$k''(\mu)$	6	0, $\tilde{C}_{[6,11]}(?)$		
12	$\kappa_2 \mapsto \nu$	k''			1		

$$[*] = \tilde{C}_{[1,9]}(\tilde{C}_{[1,6]}(1) - 8) - 8$$

TABLE 1. Data for the construction of $\tilde{a}_m^{(n)}$.

umn 6 to produce $\tilde{C}_{[1,6]}(1)$ embeddings in column 1, with $\tilde{\alpha}_8^{(1)} = \mu$ and $\tilde{\alpha}_{\tilde{C}_{[1,6]}(1)}^{(1)} = \nu$. Apply $\tilde{a}_0^{(10)}$ to all but the first 8 of these in order to get the first $\tilde{C}_{[1,6]}(1) - 8$ embeddings in column 9, with $\tilde{\alpha}_0^{(9)} = \xi$ and $\tilde{\alpha}_{\tilde{C}_{[1,6]}(1)-8}^{(9)} = \tilde{\alpha}_0^{(10)}$. These can be used to produce $\tilde{C}_{[1,9]}(\tilde{C}_{[1,6]}(1) - 8)$ embeddings in column 1, including the ones produced before; apply $\tilde{a}_0^{(10)}$ to all but the first 8 of these in order to get the first $\tilde{C}_9(1)$ embeddings in column 9 (including the ones produced before). Now, given the $\tilde{C}_9(n)$ embeddings in column 9 produced from n embeddings in column 10 ($n \geq 1$), do the previous columns' constructions to get $\tilde{C}_{[1,9]}(\tilde{C}_9(n))$ embeddings in column 1, and apply $\tilde{a}_n^{(10)}$ to all but the first $\tilde{C}_{[1,6]}(1)$ of these to get new embeddings in column 9.

Building column 11 from column 10 requires the assumption $\tilde{a}_0^{(11)}(j)(\mu) = \xi$. The construction is straightforward: given $\tilde{C}_{10}(n)$ embeddings in column 10, process the preceding columns to get $\tilde{C}_{[3,10]}(\tilde{C}_{10}(n))$ embeddings in column 3, and apply $\tilde{a}_n^{(11)}$ to all of these to get new embeddings for column 10.

Finally, in building column 11, one does not have a true column 12 but just a single embedding k'' . One gets column 11 by applying k'' to column 6: $\tilde{a}_i^{(11)} = k''(\tilde{a}_i^{(6)})$ and $\tilde{\alpha}_i^{(11)} = k''(\tilde{\alpha}_i^{(6)})$ for all i . Since $\tilde{C}_{[6,11]}(n) > n$ for all n , there is no problem continuing this indefinitely to make column 11 as long as desired.

This entire construction is summarized in Table 1, which includes: the beginning of the critical sequence of $\tilde{a}_m^{(n)}$; the initial boxed entry in column n , if any; the value of $\tilde{\alpha}_0^{(n)}$, usually used as an initial condition in the construction; the number $\hat{n}$ such that column n is extended by applying embeddings from column $n+1$ to embeddings from column $\hat{n}$; the initial value for $\tilde{C}_n$, or the initial two values if necessary; the recursive formula for $\tilde{C}_n(m+1)$ (which does not apply to $m=0$ if two values were given in the preceding column); and an estimate for the growth rate of $\tilde{C}_n(m)$.

Straightforward inductions show that the values given under “Growth” are actually lower bounds on $\tilde{C}_n(m)$. In fact, we have $\tilde{C}_n(m) = F_3(m-3) + 3$ in the columns where $\tilde{C}_n(m) = 2^m$, and $\tilde{C}_n(m) \geq F_i(m-d)+4$ in the other columns, for the listed values of i and d ; this trivial improvement makes the inductions easier, and also makes it easier to compose the estimates in the following calculations.

In order to conclude anything about $F(4)$ from this construction, we need to know that certain critical points lie below κ_4 . For example, once we know that $\xi < \kappa_4$, we can conclude that $F(4) > \tilde{C}_{[0,8]}(1)$; the estimates in Table 1 allow us to compute $\tilde{C}_{[0,8]}(1) = \tilde{C}_{[0,7]}(\tilde{C}_{[1,6]}(1) - 8) > F_6(F_4(F_4(254)))$. A better result can be obtained from the inequality $k''(k''(\mu)) < \kappa_4$. Since $k''(\mu) = k''(k(\kappa_2)) = k''(k)(k''(\kappa_2)) = \tilde{a}_0^{(11)}(\nu) = \tilde{\alpha}_0^{(11)} = \tilde{\alpha}_{\tilde{C}_{[3,6]}(1)}^{(10)}$, we get $\tilde{C}_{[6,10]}(\tilde{C}_{[3,6]}(1))$ embeddings in column 6 before reaching the ordinal $k''(\mu)$. Therefore, we get $\tilde{C}_{[6,10]}(\tilde{C}_{[3,6]}(1))$ embeddings in column 11 before reaching $k''(k''(\mu))$, so $F(4) > \tilde{C}_{[0,11]}(\tilde{C}_{[6,10]}(\tilde{C}_{[3,6]}(1))) > F_9(F_8(F_4(254)))$.

Finally, by pushing the computations in the next section a little farther, we will show that $k''(k''(k)(k''(k)(k(j)(\mu)))) < \kappa_4$. Since $k(j)(\mu) = \tilde{\alpha}_1^{(5)}$, we get $\tilde{C}_{[3,5]}(1)$ embeddings in column 3 before reaching $k(j)(\mu)$, and this leads to a total of $\tilde{C}_{[0,11]}(\tilde{C}_{[6,10]}(\tilde{C}_{[3,10]}(\tilde{C}_{[3,5]}(1))))$ critical points produced below $k''(k''(k)(k''(k)(k(j)(\mu))))$. This gives a lower bound of $F_9(F_8(F_8(254)))$ for $F(4)$. One could pursue the inequalities further, but we have clearly reached the point of diminishing returns.

There is no reason whatsoever to believe that the results in this section are even close to optimal; the construction here is merely the most successful from a series of *ad hoc* attempts to modify the original construction to produce critical points below κ_4 . It is probable that further modifications of the scheme would improve the lower bound on $F(4)$. One idea for modification would be to try to produce a true column 12 instead of a single embedding, and it would appear that the critical sequence for a member of this column should begin with $\kappa_2 \mapsto \nu \mapsto \tilde{\alpha}_m^{(12)} \mapsto \tilde{\alpha}_{m+1}^{(12)}$. Unfortunately, this will not work with the present k'' , because it turns out that $k''(k''(\nu))$ is greater than κ_4 . So apparently more drastic revisions would be needed.

5. CRITICAL POINT INEQUALITIES

In this section, we will use several basic methods to perform a number of computations on critical points. The main goal is to produce embeddings k , k' , and k'' satisfying the assumptions stated in section 4, but the accumulated facts may be useful for other purposes (e.g., as a place to search for improved versions of the construction in section 4).

We will be concentrating on the sequence $j_{(n)}$ defined earlier by $j_{(1)} = j$ and $j_{(n+1)} = j_{(n)}(j)$. When looking at these embeddings and their combinations, it will be helpful to omit some of the parentheses in applications, thus writing jj instead of $j(j)$. In larger combinations, applications will be associated from the left: $e_1e_2e_3$ means $(e_1e_2)e_3$, which is short for $(e_1(e_2))(e_3)$. So one could write $jjjj$ for $j_{(4)}$.

A great deal about the critical points of these embeddings can be deduced from the simple rules $cr(e_1e_2) = e_1(cr(e_2))$ and $e_1e_2(e_1(\beta)) = e_1(e_2(\beta))$. For example, this sufficed in section 1 to show

that $j_{(3)}(\kappa_1)$ is strictly between κ_2 and κ_3 . (Since this ordinal will arise frequently, it will be useful to have a name for it; let us call it $\kappa_{2.5}$.) Furthermore, all of the critical point manipulations in sections 2 through 4 (except for the assumptions in section 4 which were explicitly postponed until this section) used only these two rules.

In order to proceed much further with the sequence $j_{(n)}$, though, we need to use additional methods. One very useful method is to explicitly use the fact that $e_1 e_2$ is obtained by applying e_1 to initial segments of e_2 : $e_1 e_2 \upharpoonright V_{e_1(\alpha)} = e_1(e_2 \upharpoonright V_\alpha)$. Hence, if e_2 agrees with e'_2 up to α (i.e., $e_2 \upharpoonright V_\alpha = e'_2 \upharpoonright V_\alpha$), then $e_1 e_2$ agrees with $e_1 e'_2$ up to $e_1(\alpha)$. Furthermore, if e_1 agrees with e'_1 up to some ordinal greater than the rank of $e_2 \upharpoonright V_\alpha$, then $e_1 e_2$ agrees with $e'_1 e'_2$ up to $e_1(\alpha)$.

Laver [5] has defined a variant form of ‘restriction’ that turns out to be more useful for these computations than the standard definition. Let $e \bigcap^* V_\beta = \{(x, y) \in V_\beta \times V_\beta : y \in e(x)\}$. We again have $e_1 e_2 \bigcap^* V_{e_1(\beta)} = e_1(e_2 \bigcap^* V_\beta)$. We can now define the modified version of ‘agreement up to β ’: say that $e \stackrel{\beta}{=} e'$ if $e \bigcap^* V_\beta = e' \bigcap^* V_\beta$. Then, for any limit ordinal β , $\stackrel{\beta}{=}$ turns out to be an equivalence relation that respects composition and application. (To prove this, suppose that $e_1 \stackrel{\beta}{=} e'_1$ and $e_2 \stackrel{\beta}{=} e'_2$, and let $x, y \in V_\beta$; fix an ordinal γ greater than the ranks of x and y but less than β . If $y \in (e_1 \circ e_2)(x) = e_1(e_2(x))$, then $y \in e_1(z)$, where $z = e_2(x) \cap V_\gamma$. Since $e_1 \stackrel{\beta}{=} e'_1$, we have $y \in e'_1(z)$; since $e_2 \stackrel{\beta}{=} e'_2$, we have $z = e'_2(x) \cap V_\gamma$. Therefore, $y \in (e'_1 \circ e'_2)(x)$. For application, if $y \in e_1 e_2(x)$, then $(x, y) \in e_1(z)$, where $z = e_2 \bigcap^* V_\gamma$. Now $e_1 \stackrel{\beta}{=} e'_1$ gives $(x, y) \in e'_1(z)$, and $e_2 \stackrel{\beta}{=} e'_2$ gives $z = e'_2 \bigcap^* V_\gamma$, so $y \in e'_1 e'_2(x)$.)

The basic facts we will use about the relation $\stackrel{\beta}{=}$ are:

- (1) $\stackrel{\beta}{=}$ is an equivalence relation;
- (2) if $e_1 \stackrel{\beta}{=} e'_1$ and $e_2 \stackrel{\beta}{=} e'_2$, then $e_1(e_2) \stackrel{\beta}{=} e'_1(e'_2)$ and $e_1 \circ e_2 \stackrel{\beta}{=} e'_1 \circ e'_2$;
- (3) if $e_2 \stackrel{\beta}{=} e'_2$, then $e_1(e_2) \stackrel{e_1(\beta)}{=} e_1(e'_2)$;
- (4) $e \stackrel{cr(e)}{=} \text{id}$, where id is the identity function; and
- (5) if $e \stackrel{\beta}{=} e'$ and either $e(\alpha)$ or $e'(\alpha)$ is less than β , then $e(\alpha) = e'(\alpha)$.

The first two of these were proved above, and the third is a consequence of the elementarity of e_1 ; the last two are easy from the definitions. [For (5), we cannot have, for example, $e(\alpha) < \beta$ and $e(\alpha) < e'(\alpha)$, because then the pair $(\alpha, e(\alpha))$ would be in $e' \bigcap^* V_\beta$ but not in $e \bigcap^* V_\beta$.]

One application of these restriction methods is the following lemma.

Lemma 15. *If $cr(e) \geq \kappa_2$, then $ejjj(\kappa_1) = e(\kappa_{2.5})$.*

Proof 1. Note that, if $e(x) = x$, then $ee'(x) = ee'(e(x)) = e(e'(x))$. Now, since κ_2 is regular, the map $j \upharpoonright \kappa_1$ from κ_1 to κ_2 is not cofinal in κ_2 , so the rank of $j \upharpoonright V_{\kappa_1}$ is less than κ_2 . Since $ej(\kappa_1) = e(j(\kappa_1)) = e(\kappa_2) \geq \kappa_2$, κ_1 and $j \upharpoonright \kappa_1$ are in the domain of $ej(j \upharpoonright V_{\kappa_1})$, and we have $ej(j \upharpoonright V_{\kappa_1})(\kappa_1) = e(j(j \upharpoonright V_{\kappa_1})(\kappa_1)) = e(\kappa_2) > \kappa_1$. Therefore,

$$ejjj(\kappa_1) = ej(j \upharpoonright V_{\kappa_1})(j \upharpoonright V_{\kappa_1})(\kappa_1) = e(j(j \upharpoonright V_{\kappa_1})(j \upharpoonright V_{\kappa_1})(\kappa_1)) = e(jjj(\kappa_1)) = e(\kappa_{2.5}).$$

Proof 2. Since $cr(e) \geq \kappa_2$, we have $e \stackrel{\kappa_2}{=} \text{id}$. This gives $ej \stackrel{\kappa_2}{=} \text{id}(j) = j$ and hence $ejj \stackrel{\kappa_2}{=} jj$, so

$$ej(\kappa_2) = ej(j(\kappa_1)) = ejj(ej(\kappa_1)) \geq ejj(\kappa_2) = ejjj(ejj(\kappa_1)) \geq ejjj(\kappa_2) > ejjj(\kappa_1).$$

We now have $ejjj \stackrel{ejj(\kappa_2)}{=} ejj(ej) \stackrel{ej(\kappa_2)}{=} ej(ej)(ej) = e(jjj)$. Therefore, $ejjj(\kappa_1) = e(jjj)(\kappa_1) = e(jjj)(e(\kappa_1)) = e(jjj(\kappa_1)) = e(\kappa_{2.5})$. ■

From now on, the derivations in this section will be in the form of Proof 2 (using modified restrictions) rather than that of Proof 1 (using standard restrictions). Derivations using standard restrictions are sometimes easier to come up with—the above lemma is an example—but derivations using modified restrictions give a little more information and are applicable in other contexts besides that of elementary embeddings (as in forthcoming work of Jech and myself [3]).

Using this lemma and the basic rules, we can now derive more information about the embeddings $j_{(n)}$ for relatively small n , as shown in Table 2. In this table and from now on, κ_i^n is an abbreviation for $j_{(n)}(\kappa_i)$.

n	Critical sequence of $j_{(n)}$	Other values	n	Critical sequence of $j_{(n)}$	Other values
1	$\kappa_0 \mapsto \kappa_1 \mapsto \kappa_2 \mapsto \kappa_3 \mapsto \kappa_4$		9	$\kappa_0 \mapsto \kappa_1 \mapsto \kappa_2 \mapsto \kappa_2^7 \mapsto >\kappa_4$	
2	$\kappa_1 \mapsto \kappa_2 \mapsto \kappa_3 \mapsto \kappa_4$		10	$\kappa_1 \mapsto \kappa_2 \mapsto \kappa_2^7 \mapsto \kappa_3^9 \mapsto >\kappa_4$	
3	$\kappa_0 \mapsto \kappa_2 \mapsto \kappa_3 \mapsto \kappa_4$	$\kappa_1 \mapsto \kappa_{2.5}$	11	$\kappa_0 \mapsto \kappa_2 \mapsto \kappa_2^7 \mapsto \kappa_3^{10} \mapsto >\kappa_4$	$\kappa_1 \mapsto \kappa_3$
4	$\kappa_2 \mapsto \kappa_{2.5} \mapsto \kappa_3 \mapsto \kappa_4$		12	$\kappa_2 \mapsto \kappa_3 \mapsto \kappa_2^7 \mapsto \kappa_3^{11} \mapsto >\kappa_4$	
5	$\kappa_0 \mapsto \kappa_1 \mapsto \kappa_{2.5} \mapsto \kappa_4$		13	$\kappa_0 \mapsto \kappa_1 \mapsto \kappa_3 \mapsto \kappa_2^7 \mapsto >\kappa_4$	
6	$\kappa_1 \mapsto \kappa_{2.5} \mapsto \kappa_2^5 \mapsto >\kappa_4$		14	$\kappa_1 \mapsto \kappa_3 \mapsto \kappa_2^{13} \mapsto \kappa_2^7 \mapsto >\kappa_4$	
7	$\kappa_0 \mapsto \kappa_{2.5} \mapsto \kappa_2^6 \mapsto \kappa_3^6 \mapsto >\kappa_4$	$\kappa_1 \mapsto \kappa_3$	15	$\kappa_0 \mapsto \kappa_3 \mapsto \kappa_2^{14} \mapsto \kappa_2^{13} \mapsto >\kappa_4$	
8	$\kappa_{2.5} \mapsto \kappa_3 \mapsto \kappa_2^7 \mapsto \kappa_3^7 \mapsto >\kappa_4$		16	$\kappa_3 \mapsto \kappa_1^{15} \mapsto \kappa_2^{15} \mapsto \kappa_2^{14} \mapsto >\kappa_4$	

TABLE 2. Selected values of $j_{(n)}$ for $1 \leq n \leq 16$.

The ‘ $>\kappa_4$ ’ entry for $j_{(6)}$ comes from $\kappa_3^5 > \kappa_{2.5}^5 = \kappa_4$. For the ‘ $>\kappa_4$ ’ entry for $j_{(n)}$ ($7 \leq n \leq 16$), we need the fact that $j_{(n-1)}(\kappa_4) > \kappa_4$. An easy way to see this is to prove by induction that an ordinal fixed by an embedding e which lies above the critical point of e must lie above the entire critical sequence of e . Another way is to conclude from a theorem of Kunen (in the form given in Solovay-Reinhardt-Kanamori [8]) or the results of Laver and Steel mentioned previously that the critical sequence of e must be cofinal in λ . (Recall that $j: V_\lambda \rightarrow V_\lambda$.)

Since Laver [6] has shown that there are exactly m critical points below $cr(j_{(2^m)})$ for all m , the fact that $\kappa_3 = cr(j_{(16)})$ implies that there are no critical points below κ_3 other than the four we already know about.

We now have enough information to prove a number of the facts needed for section 4. Let $k = j_{(10)}$; then k has critical sequence beginning $\kappa_1 \mapsto \kappa_2 \mapsto \mu \mapsto \nu$, where $\mu = \kappa_2^7$ and $\nu = \kappa_3^9$. Also, $k(j)(\kappa_1) = j_{(11)}(\kappa_1) = \kappa_3$. Next, let $k' = j_{(10)}(j_{(11)})$; k' has critical sequence beginning $j_{(10)}(\kappa_0) \mapsto j_{(10)}(\kappa_2) \mapsto j_{(10)}(\kappa_2^7) \mapsto j_{(10)}(\kappa_3^{10})$, which is $\kappa_0 \mapsto \mu \mapsto \nu \mapsto \xi$, where $\xi = j_{(10)}(\kappa_3^{10})$. Finally, if we let $k'' = j_{(9)}(j_{(14)})$, then the critical sequence of k'' will begin with $j_{(9)}(\kappa_1) \mapsto j_{(9)}(\kappa_3)$, which is $\kappa_2 \mapsto \nu$. All that remains is to show that $k''(k)(j)(\mu) = \xi$ and that the relevant critical points lie below κ_4 .

The main problem we will have to solve is that of comparing various critical points. Certain such comparisons can be determined directly from Table 2; for example, $\kappa_2^7 < \kappa_2^6$ because $\kappa_2^6 = \kappa_{2.5}^7$. A

thorough examination of Table 2 gives the following list of critical points of the forms κ_i and κ_i^n :

$$\begin{aligned} \kappa_0 < \kappa_1 < \kappa_2 < \kappa_{2.5} < \kappa_3 < \kappa_1^{15} < \kappa_2^{15} < \kappa_{2.5}^{15} < \kappa_2^{14} < \kappa_{2.5}^{14} < \kappa_2^{13} < \kappa_{2.5}^{13} \\ < \kappa_2^7 < \kappa_2^6 < \kappa_2^5 < \kappa_4 < \kappa_3^5. \end{aligned}$$

However, the ordinals $\kappa_{2.5}^n$ for $n = 9, 10, 11$ and κ_3^n for $n = 6, 7, 9, 10, 11$ cannot yet be placed in this list; we only have the partial information that $\kappa_2^7 < \kappa_{2.5}^{11} < \kappa_3^{11} < \kappa_3^{10}$, $\kappa_2^7 < \kappa_{2.5}^{10} < \kappa_3^{10} < \kappa_3^9$, $\kappa_2^7 < \kappa_{2.5}^9 < \kappa_3^9$, $\kappa_2^6 < \kappa_3^7 < \kappa_3^6$, and $\kappa_2^5 < \kappa_3^6$. The critical points $\kappa_{2.5}^n$ for $n = 1, 2, 3, 12$ are not mentioned here because, as we will see soon, they already occur in the above list. We will return later to the problem of sorting the remaining critical points into the correct order.

One useful method for critical point computations is to approximate (in the sense of $\overset{\alpha}{\equiv}$) complicated expressions such as $j_{(n)}$ for relatively large n by simpler expressions. In order to do this, some basic computations are useful.

First, from Laver [5], for any embeddings e and $e_1, \dots, e_l$, we have $ee_1e_2 \dots e_l \overset{\theta}{=} e(e_1e_2 \dots e_l)$, where θ is the minimum of $ee_1e_2 \dots e_i(cr(e))$ for $1 \leq i < l$. This is proved by induction:

$$ee_1e_2 \dots e_l \overset{ee_1e_2 \dots e_{l-1}(cr(e))}{=} ee_1e_2 \dots e_{l-1}(ee_l) \overset{\theta}{=} e(e_1e_2 \dots e_{l-1})(ee_l) = e(e_1e_2 \dots e_l).$$

Second, we have $e \overset{e(cr(e'))}{=} e \circ e'$. To see this, note that $cr(ee') = e(cr(e'))$, so $\text{id} \overset{e(cr(e'))}{=} ee'$, so $e = \text{id} \circ e \overset{e(cr(e'))}{=} ee' \circ e = e \circ e'$. A special case of this is $ee' \overset{ee'(cr(e))}{=} ee' \circ e = e \circ e'$. This is a version of the fact given earlier, that $e(x) = x$ implies $ee'(x) = ee'(e(x)) = e(e'(x))$.

Using these facts, we can see that $j_{(3)}$ is approximated by $j \circ j$, because $j_{(3)} \overset{\kappa_{2.5}}{=} j_{(2)} \circ j = j \circ j$. But a glance at Table 2 indicates that $j \circ j$ should approximate $j_{(11)}$ better than it approximates $j_{(3)}$, since $j_{(11)}(\kappa_1) = (j \circ j)(\kappa_1) = \kappa_3 \neq j_{(3)}(\kappa_1)$. To see that this is indeed the case, note that $j_{(11)} = j_{(8)}jjj \overset{\kappa_2^7}{=} j_{(8)}(jjj) = j_{(8)}j_{(3)}$, since both $j_{(8)}j(\kappa_{2.5})$ and $j_{(8)}jj(\kappa_{2.5})$ are greater than κ_2^7 . Similarly, we get $j_{(8)} \overset{\kappa_2^7}{=} j_{(4)}j_{(4)} = j_{(3)}j_{(2)}$, so $j_{(11)} \overset{\kappa_2^7}{=} j_{(3)}j_{(2)}j_{(3)}$. Now note that

$$j_{(3)}j_{(2)}j_{(3)}(\kappa_{2.5}) > j_{(3)}j_{(2)}j_{(3)}(\kappa_2) = j_{(3)}j_{(2)}(j_{(3)}(\kappa_2)) = j_{(3)}(\kappa_3) = \kappa_4 > \kappa_2^7,$$

so $j_{(11)} \overset{\kappa_2^7}{=} j_{(3)}j_{(2)}j_{(3)} \overset{\kappa_2^7}{=} j_{(3)}j_{(2)} \circ j_{(3)} = j_{(3)} \circ j_{(2)} = j_{(2)} \circ j = j \circ j$. Since $j_{(11)}(\kappa_2) = \kappa_2^7$, we get as a consequence that $j_{(11)} \upharpoonright V_{\kappa_2} = (j \circ j) \upharpoonright V_{\kappa_2}$.

Similarly, one can approximate $j_{(12)}$ by $jj_{(2)}$, since $j_{(12)} = j_{(11)}j \overset{\kappa_2^7}{=} (j \circ j)j = jj_{(2)}$. Continuing, we get the following:

$$\begin{aligned} j_{(13)} &= j_{(12)}j \overset{\kappa_2^7}{=} jj_{(2)}j \\ j_{(14)} &= j_{(12)}jj \overset{\kappa_{2.5}^{13}}{=} j_{(12)}(jj) \overset{\kappa_2^7}{=} jj_{(2)}(jj) = jj_{(3)} \\ j_{(15)} &= j_{(12)}jjj \overset{\kappa_2^{14}}{=} j_{(12)}(jjj) \overset{\kappa_2^7}{=} jj_{(2)}(jjj) = j_{(2)}j_{(2)}(j_{(2)}j) = j_{(2)}j_{(3)}. \end{aligned}$$

These approximations allow us to evaluate certain critical points in terms of others. First, since $j_{(14)}(\kappa_2) < \kappa_2^{13}$, we have $\kappa_2^{14} = jj_{(3)}(\kappa_2) = jj_{(3)}(j(\kappa_1)) = j(j_{(3)}(\kappa_1)) = \kappa_{2.5}^1$. Similarly, $\kappa_2^{15} < \kappa_2^{14}$ implies $\kappa_2^{15} = j_{(2)}j_{(3)}(\kappa_2) = j_{(2)}j_{(3)}(j_{(2)}(\kappa_1)) = \kappa_{2.5}^2$. Since $\kappa_1^{15} < \kappa_2^{14}$, we have $\kappa_1^{15} =$

$j_{(12)}(jjj)(\kappa_1) = j_{(12)}j_{(3)}(j_{(12)}(\kappa_1)) = \kappa_{2.5}^{12}$. But we can go further here; since $j_{(12)} \stackrel{\kappa_2^7}{=} jj_{(2)} = j_{(2)}j_{(2)} = j_{(3)}j_{(3)}$, we get $\kappa_{2.5}^{12} = j_{(3)}j_{(3)}(\kappa_{2.5}) = j_{(3)}j_{(3)}(j_{(3)}(\kappa_1)) = \kappa_{2.5}^3$. So the critical points $\kappa_{2.5}^n$ for $n = 1, 2, 3, 12$ are already in our list and do not need to be added.

We can now get the approximations $j_{(13)} \stackrel{\kappa_2^{13}}{=} j \circ j_{(2)}$ and $j_{(14)} \stackrel{\kappa_2^{14}}{=} j_{(2)} \circ j_{(2)}$ as follows:

$$\begin{aligned} j_{(13)} &\stackrel{j_{(13)}(cr(jj_{(2)}))}{=} j_{(13)} \circ jj_{(2)} \stackrel{\kappa_2^7}{=} jj_{(2)}j \circ jj_{(2)} = jj_{(2)} \circ j = j \circ j_{(2)}, \\ j_{(14)} &\stackrel{\kappa_2^{13}}{=} jj_{(3)} \stackrel{j(\kappa_{2.5})}{=} j(j \circ j) = j_{(2)} \circ j_{(2)}. \end{aligned}$$

The latter implies that $j_{(15)} \stackrel{\kappa_1^{15}}{=} j \circ j \circ j$, since

$$j_{(15)} = j_{(14)}j \stackrel{\kappa_1^{15}}{=} j_{(14)} \circ j \stackrel{\kappa_2^{14}}{=} j_{(2)} \circ j_{(2)} \circ j = j_{(2)} \circ j \circ j = j \circ j \circ j.$$

This is not as good an approximation for $j_{(15)}$ as $j_{(2)}j_{(3)}$, but it will suffice for the following.

We are now ready to show that $k''(k)(j)(\mu) = \xi$. Since $k'' = j_{(9)}j_{(14)}$ and $k = j_{(10)} = j_{(9)}j$, we have $k''k = j_{(9)}j_{(15)}$. We also have $j_{(9)}j_{(15)} \stackrel{j_{(9)}(\kappa_1^{15})}{=} j_{(9)}(j \circ j \circ j)$; since

$$\begin{aligned} j_{(9)}(j_{(15)}(\kappa_1)) &= j_{(9)}j_{(15)}(j_{(9)}(\kappa_1)) = j_{(9)}j_{(15)}(\kappa_2) = j_{(9)}j_{(15)}(j(\kappa_1)) = j_{(9)}j_{(15)}j(j_{(9)}j_{(15)}(\kappa_1)) \\ &= j_{(9)}j_{(15)}j(j_{(9)}j_{(15)}(j_{(9)}(\kappa_0))) = j_{(9)}j_{(15)}j(\kappa_3^9) > j_{(9)}j_{(15)}j(\mu), \end{aligned}$$

we get

$$\begin{aligned} j_{(9)}j_{(15)}j(\mu) &= j_{(9)}(j \circ j \circ j)j(\mu) = (j_{(10)} \circ j_{(10)} \circ j_{(10)})j(\mu) \\ &= j_{(10)}(j_{(10)}(j_{(11)}))(j_{(10)}(j_{(10)}(\kappa_1))) = j_{(10)}(j_{(10)}(\kappa_3)) = \xi. \end{aligned}$$

It remains to show that a certain critical point lies below κ_4 . In order to do so, we will study a more general question: Given an embedding e and a critical point γ , what is the least α such that $e(\alpha) \geq \gamma$? (This will be solved here only in a few special cases; in general it seems quite difficult.) This ordinal α is often not a critical point, so we will start by studying some additional ordinals which are definable from the embeddings in $\mathcal{P}_j$.

For any embedding e and ordinal α , let $e(<\alpha)$ be the strict supremum of $e(\beta)$ for $\beta < \alpha$ (i.e., the least ordinal greater than all such $e(\beta)$). We clearly have $e(<\alpha) \leq e(\alpha)$. If ρ is the cofinality of α , then $e(\rho) > \rho$ implies $e(<\alpha) < e(\alpha)$, because the cofinality of $e(<\alpha)$ is ρ while that of $e(\alpha)$ is $e(\rho)$. (Conversely, if $e(\rho) = \rho$, then $e(<\alpha) = e(\alpha)$. To see this, let $g: \rho \rightarrow \alpha$ be increasing and cofinal in α ; then $e(g)$ must be increasing and cofinal in $e(\alpha)$. But the domain of $e(g)$ is $e(\rho) = \rho$, and $e(g)(e(\delta)) = e(g(\delta))$ for $\delta < \rho$, so the ordinals $e(\beta)$ for $\beta < \alpha$ are cofinal in $e(\alpha)$.)

As usual, elementarity implies that application distributes over this new operation: $e'(e(<\alpha)) = e' e(<e'(\alpha))$. Also, for any e' , e , and α , the ordinals $e(\beta)$ for $\beta < \alpha$ are cofinal in $e(<\alpha)$, so the ordinals $e'(e(\beta))$ for $\beta < \alpha$ are cofinal in $e'(<e(<\alpha))$; this gives the formula $e'(<e(<\alpha)) = (e' \circ e)(<\alpha)$. Another useful fact is that $e \stackrel{e(\alpha)}{=} e'$ implies $e(\beta) = e'(\beta)$ for $\beta < \alpha$, and hence $e(<\alpha) = e'(<\alpha)$.

Two of these ordinals arise frequently enough that it is useful to have short names for them: let $\sigma_1 = j(<\kappa_1)$ and $\sigma_2 = j(<\kappa_2)$. Then we can see from the above that σ_1 is strictly between

κ_1 and κ_2 , while σ_2 is strictly between κ_2 and κ_3 . To see where σ_2 lies relative to $\kappa_{2.5}$, use the following computation:

$$\sigma_2 = j(<\kappa_2) > j(\sigma_1) = j_{(2)}(<\kappa_2) > j_{(2)}(\sigma_1) = j_{(3)}(<\kappa_2) > j_{(3)}(\kappa_1) = \kappa_{2.5}.$$

On the other hand, the fact that $j_{(3)} \stackrel{\kappa_{2.5}}{\cong} j \circ j$ yields

$$j(<\sigma_1) = (j \circ j)(<\kappa_1) = j_{(3)}(<\kappa_1) < j_{(3)}(\kappa_1) = \kappa_{2.5}.$$

So $\kappa_{2.5}$ lies within a gap in the range of j , and σ_1 is the least ordinal that j sends above $\kappa_{2.5}$.

The inequality $(j \circ j)(<\kappa_1) < \kappa_{2.5}$ will be very useful later, because we can apply an arbitrary embedding e to it to get $(ej \circ ej)(<e(\kappa_1)) < e(\kappa_{2.5})$. In other words,

$$(\dagger) \quad \text{if } \beta < e(\kappa_1), \quad \text{then } ej(ej(\beta)) < e(\kappa_{2.5}).$$

We can now try to determine the ordinals at which various embeddings first jump beyond κ_4 . Let us first consider $j_{(6)}$; the ordinal here turns out to be $j_{(5)}(\sigma_1)$. To see this, compute as follows:

$$\begin{aligned} j_{(6)}(j_{(5)}(\sigma_1)) &= j_{(5)}(j(\sigma_1)) > j_{(5)}(\kappa_{2.5}) = \kappa_4, \\ j_{(6)}(<j_{(5)}(\sigma_1)) &= j_{(5)}(j(<\sigma_1)) < j_{(5)}(\kappa_{2.5}) = \kappa_4. \end{aligned}$$

In particular, since $j_{(5)}(\sigma_1) = j_{(6)}(<\kappa_{2.5}) > \kappa_2^6$, we get $\kappa_3^6 < j_{(6)}(\kappa_2^6) < \kappa_4$.

[One could give a proof of $j_{(6)}(\kappa_2^6) < \kappa_4$ without using ordinals other than critical points, as follows: we have $j_{(3)} \stackrel{\kappa_{2.5}}{\cong} j \circ j$, so $j_{(5)}j_{(3)} \stackrel{\kappa_4}{\cong} j_{(5)}(j \circ j)$, so

$$\kappa_4 = j_{(5)}(j_{(3)}(\kappa_1)) = j_{(5)}j_{(3)}(\kappa_{2.5}) > j_{(5)}j_{(3)}(\kappa_2) = j_{(5)}(j \circ j)(\kappa_2) = j_{(6)}(j_{(6)}(\kappa_2)).$$

In fact, Jech and I have shown [3] that, in theory, any argument involving the ordinals $e(<\alpha)$ and the methods used here can be translated into an argument referring only to critical points. However, such arguments tend to involve long, roundabout, and opaque computations.]

The embedding $j_{(9)}$ is a more difficult case. The least ordinal sent above κ_4 by $j_{(9)}$ is $j_{(8)}(\sigma_2)$. To see that $j_{(9)}(j_{(8)}(\sigma_2)) > \kappa_4$, note that

$$\begin{aligned} j(\sigma_2) &= j_{(2)}(<\kappa_3) > j_{(2)}(\sigma_2) = j_{(3)}(<\kappa_3) > j_{(3)}(\sigma_2) = j_{(4)}(<\kappa_3) > j_{(4)}(\sigma_2) = j_{(5)}(<\kappa_{2.5}) \\ &> j_{(5)}(j(<\sigma_1)) = j_{(6)}(<j_{(5)}(\sigma_1)) > \kappa_3^6 > \kappa_3^7, \end{aligned}$$

so $j_{(9)}(j_{(8)}(\sigma_2)) = j_{(8)}(j(\sigma_2)) > j_{(8)}(\kappa_3^7) > \kappa_4$. For the other side, first use the fact that $j_{(10)} \stackrel{\kappa_2^7}{\cong} j_{(8)}(j_{(2)})$ (since $\kappa_{2.5}^9 > \kappa_2^9 = \kappa_2^7$) to see that $j_{(10)}(\sigma_1) = j_{(8)}(jj)(\sigma_1) = j_{(8)}(j_{(2)}(\sigma_1))$ (since $\sigma_1 < \kappa_2$ and $\kappa_2^{10} = \kappa_2^7$). From this, we get

$$\begin{aligned} j_{(9)}(<j_{(8)}(\sigma_2)) &= j_{(8)}(j(<\sigma_2)) = j_{(8)}((j \circ j)(<\kappa_2)) = j_{(8)}(j_{(11)}(<\kappa_2)) \\ &= j_{(8)}(j_{(10)}(\sigma_1)) = j_{(8)}(j_{(8)}(j_{(2)}(\sigma_1))). \end{aligned}$$

Since $j_{(2)}(\alpha_1) < \kappa_3 = j_{(7)}(\kappa_1)$, we can now apply $(\dagger)$ to get $j_{(8)}(j_{(8)}(j_{(2)}(\sigma_1))) < j_{(7)}(\kappa_{2.5}) = \kappa_2^6 < \kappa_4$.

By methods similar to the above (with each n handled separately), it can be shown that: for $7 \leq n \leq 13$, the least α such that $j_{(n)}(\alpha) > \kappa_4$ is $j_{(n-1)}(\sigma_2)$; for $n = 14$ or $n = 15$, the least such α is $j_{(n-1)}(j(\sigma_1))$.

We now have the needed tools to complete the ordering of the critical points κ_i^n for $i \leq 3$ and $n \leq 16$. The main tool needed here is the formula $j(<\sigma_1) < \kappa_{2.5}$; as with $(\dagger)$, we can apply any embedding e to this to get $ej(<e(\sigma_1)) < e(\kappa_{2.5})$. For instance, take $e = j_{(9)}$ to get $j_{(10)}(<j_{(9)}(\sigma_1)) < \kappa_{2.5}^9$; from this and the calculation

$$j_{(9)}(\sigma_1) = j_{(10)}(<\kappa_2) > j_{(10)}(\sigma_1) = j_{(11)}(<\kappa_2) > j_{(11)}(\sigma_1) = j_{(12)}(<\kappa_3) > \kappa_2^{12} = \kappa_3,$$

we get $\kappa_3^{10} < \kappa_{2.5}^9$. The same argument shows that $\kappa_3^{11} < \kappa_{2.5}^{10}$; it also shows that $\kappa_3^7 < \kappa_{2.5}^6 = \kappa_2^5$, once we verify that $j_{(6)}(\sigma_1) > \kappa_3$. To see this, note that $j_{(4)}jj \stackrel{\kappa_2^5}{=} j_{(4)}(jj)$ and $\kappa_2^5 > \kappa_2^6 > j_{(6)}(\sigma_1)$, so $j_{(6)}(\sigma_1) = j_{(4)}j_{(2)}(\sigma_1) = j_{(4)}(j_{(2)}(\sigma_1)) > j_{(4)}(\kappa_{2.5}) = \kappa_3$.

It remains to compare κ_3^9 with κ_2^6 ; we can show that $\kappa_3^9 < \kappa_2^6$ as follows. Apply $j_{(7)}$ to the formula $j(<\sigma_1) < \kappa_{2.5}$ to get $j_{(8)}(<j_{(7)}(\sigma_1)) < \kappa_{2.5}^7 = \kappa_2^6$. Since $j_{(9)}(\kappa_3) = j_{(9)}(j_{(8)}(\kappa_{2.5})) = j_{(8)}(j(\kappa_{2.5}))$, all we need to show is that $j(\kappa_{2.5}) < j_{(7)}(\sigma_1)$. This is proved by the following computation, part of which was already done in the preceding paragraph:

$$\begin{aligned} j_{(7)}(\sigma_1) &= j_{(8)}(<\kappa_3) > j_{(8)}(\sigma_2) = j_{(9)}(<\kappa_2) > j_{(9)}(\sigma_1) \\ &> j_{(12)}(<\kappa_3) > j_{(12)}(\sigma_2) = j_{(13)}(<\kappa_3) > \kappa_{2.5}^{13} > \kappa_2^{14} = \kappa_{2.5}^1. \end{aligned}$$

This completes the comparison of these critical points κ_i^n ; the part of the list from κ_2^7 on is

$$\kappa_2^7 < \kappa_{2.5}^{11} < \kappa_3^{11} < \kappa_{2.5}^{10} < \kappa_3^{10} < \kappa_{2.5}^9 < \kappa_3^9 < \kappa_2^6 < \kappa_3^7 < \kappa_2^5 < \kappa_3^6 < \kappa_4 < \kappa_3^5.$$

We are finally ready to prove the last assumption in section 4, $k''(k''(k)(k''(k)(k(j)(\mu)))) < \kappa_4$. We have $k'' = j_{(9)}j_{(14)}$, $k''(k) = j_{(9)}j_{(15)}$, and $k(j) = j_{(11)}$. This gives $k(j)(\mu) = j_{(11)}(j_{(10)}(\kappa_2)) = j_{(10)}(j(\kappa_2)) = \kappa_3^{10}$. We saw above that $\kappa_3^{10} < \kappa_{2.5}^9$. Since $j_{(14)}(\kappa_1) = \kappa_3 > \kappa_{2.5}$, $(\dagger)$ implies that $j_{(15)}(j_{(15)}(\kappa_{2.5})) < j_{(14)}(\kappa_{2.5})$; we can apply $j_{(9)}$ to this to get $j_{(9)}j_{(15)}(j_{(9)}j_{(15)}(\kappa_{2.5}^9)) < j_{(9)}(\kappa_{2.5}^{14})$. In the same way, since $j_{(13)}(\kappa_1) > \kappa_{2.5}$, $(\dagger)$ yields $j_{(14)}(j_{(14)}(\kappa_{2.5})) < j_{(13)}(\kappa_{2.5})$ and hence $j_{(9)}j_{(14)}(j_{(9)}(\kappa_{2.5}^{14})) < j_{(9)}(\kappa_{2.5}^{13})$. We saw in the preceding paragraph that $j_{(8)}(\sigma_2) > \kappa_{2.5}^{13}$, so the analysis of $j_{(9)}$ done earlier implies that $j_{(9)}(\kappa_{2.5}^{13}) < \kappa_4$. Putting all of this together, we get

$$\begin{aligned} k''(k''(k)(k''(k)(k(j)(\mu)))) &= j_{(9)}j_{(14)}(j_{(9)}j_{(15)}(j_{(9)}j_{(15)}(\kappa_3^{10}))) \\ &< j_{(9)}j_{(14)}(j_{(9)}j_{(15)}(j_{(9)}j_{(15)}(\kappa_{2.5}^9))) \\ &< j_{(9)}j_{(14)}(j_{(9)}(\kappa_{2.5}^{14})) \\ &< j_{(9)}(\kappa_{2.5}^{13}) \\ &< \kappa_4, \end{aligned}$$

as desired.

The comparisons between critical points in this section are rather haphazard; it would be nice to have a definite algorithm for deciding which of two given critical points is smaller. Such an algorithm is known [6], but it proceeds through all smaller critical points, and hence is useless if one wants to know whether, say, $j_{(12)}(j_{(13)}(\kappa_2^{13}))$ lies below $\kappa_{2.5}^{10}$. A definite question one can ask is

whether there is a primitive recursive algorithm for comparing the critical points of two embeddings (given as expressions in j).

The ordinals $e(<\alpha)$ also yield some interesting questions. For instance, the fact that there are only finitely many modified restrictions of members of $\mathcal{P}_j$ to a given critical point [6] means that there can only be finitely many such ordinals between two successive critical points; in fact, there are at most $n2^n$ of them below critical point number n if the critical points are numbered in increasing order starting with $n = 0$. But this bound is not sharp; it turns out that there are no such ordinals between κ_0 and κ_1 , one (σ_1) between κ_1 and κ_2 , one ($j_{(3)}(<\kappa_1)$) between κ_2 and $\kappa_{2.5}$, and five ($j_{(7)}(<\kappa_1)$, $j_{(3)}(\sigma_1)$, $j_{(2)}(\sigma_1)$, $j(\sigma_1)$, and σ_2) between $\kappa_{2.5}$ and κ_3 . One can ask for improved bounds on the number of these ordinals.

A more interesting question is whether the place at which an embedding e skips over a critical point β (i.e., the least α such that $e(\alpha) \geq \beta$) must be of the form $e'(<\gamma)$ for some embedding e' and critical point γ . The converse turns out to be true: if e is an embedding with critical point γ , then $e'(<\gamma)$ is the least α such that $e(\alpha) \geq ee'(\gamma)$.

6. CONCLUSION

It seems likely that the results in this paper are not optimal. However, pushing the lower bound on the growth rate of the number $F(n)$ of critical points below κ_n to a function beyond $F_{\omega+1}$ will probably require a new idea, just as it took an additional idea to get from F_ω to $F_{\omega+1}$. One might hope that the growth rate can be shown to be so great that the function F cannot be proven to be well-defined within Peano arithmetic, or perhaps even a stronger theory. In fact, one might hope to obtain the consistency of large cardinals from the assumption that $F(n)$ is finite for all n , but this would seem to be a great deal to ask for.

The assumptions on which the construction in section 4 was based appear to be highly coincidental, but this coincidence merely allowed us to get a couple of levels higher in the F_n hierarchy than we would have without it. Further experimentation with these ordinals would quite possibly reveal more such relations, allowing for an improved construction giving even more critical points below κ_4 .

The best that one could hope for in this direction would be a version of Figure 1 in which every critical point from the algebra $\mathcal{P}_j$ would be produced by applying Lemma 2 to column 1. As a starting point, Laver [6] has shown that the ordinals $\gamma_m = cr(j_{(2^m)})$ are in fact all of the critical points from $\mathcal{P}_j$, and that $j_{(2^m)}(\gamma_m) = \gamma_{m+1}$. Results I have obtained recently (a number of which were obtained independently by A. Drápal) show that the embeddings $j_{(2^{2^m}-1)}$ have the desired form for column 1 in Figure 1: $j_{(2^{2^m}-1)}$ has critical sequence beginning $\gamma_0 \mapsto \gamma_{2^m} \mapsto \gamma_{2^{m+1}}$ for all m . There is also a natural candidate for column 2: $j_{(2^{2^m}-2)}$ has critical sequence beginning $\gamma_1 \mapsto \gamma_{2^m} \mapsto \gamma_{2^{m+1}}$. These calculations also show that the first 256 critical points from $\mathcal{P}_j$ are those produced in section 2 below κ_4 , and that the next critical point beyond these is the ordinal $\mu = j_{(7)}(\kappa_2)$ used in sections 4 and 5. This leads to some hope that the ultimate construction is indeed possible. In order to produce a growth rate beyond F_ω , though, this construction would probably require an additional recursive level to produce the starting entries in whichever columns would need them, so one can expect it to be very complex.

REFERENCES

1. P. Dehornoy, *Braid groups and left distributive operations*, Rapport de recherche 1992-1, Groupe de recherche algorithmique & logique, Université de Caen.
2. ———, *Free distributive groupoids*, J. Pure Appl. Algebra **61** (1989), 123–146.
3. R. Dougherty and T. Jech, *Finite left-distributive algebras and embedding algebras*, preprint.

4. D. Larue, *On braid words and irreflexivity*, preprint.
5. R. Laver, *The left distributive law and the freeness of an algebra of elementary embeddings*, Adv. Math. **91** (1992), 209–231.
6. ———, *On the algebra of elementary embeddings of a rank into itself*, preprint.
7. H. Rose, *Subrecursion: functions and hierarchies*, Oxford Univ. Press, Oxford, 1984.
8. R. Solovay, W. Reinhardt, and A. Kanamori, *Strong axioms of infinity and elementary embeddings*, Ann. Math. Logic **13** (1978), 73–116.

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, COLUMBUS, OH 43210